

Calculus M211

Jörg Endrullis

Indiana University Bloomington

2013

Calculating Limits using Limit Laws

We have seen that calculating limits with a calculator sometimes leads to incorrect results.

We will now see how to compute limits using **Limit Laws**:


Let c be a constant, and let $\lim_{x \rightarrow a} f(x)$ and $\lim_{x \rightarrow a} g(x)$ exist. Then

1. $\lim_{x \rightarrow a} [f(x) + g(x)] = \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x)$
2. $\lim_{x \rightarrow a} [f(x) - g(x)] = \lim_{x \rightarrow a} f(x) - \lim_{x \rightarrow a} g(x)$
3. $\lim_{x \rightarrow a} [c \cdot f(x)] = c \cdot \lim_{x \rightarrow a} f(x)$
4. $\lim_{x \rightarrow a} [f(x) \cdot g(x)] = \lim_{x \rightarrow a} f(x) \cdot \lim_{x \rightarrow a} g(x)$
5. $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)}$ if $\lim_{x \rightarrow a} g(x) \neq 0$

These laws also work for one-sided limits $\lim_{x \rightarrow a^\pm}$.

Calculating Limits using Limit Laws

- $\lim_{x \rightarrow a} [f(x) + g(x)] = \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x)$
- $\lim_{x \rightarrow a} [f(x) - g(x)] = \lim_{x \rightarrow a} f(x) - \lim_{x \rightarrow a} g(x)$
- $\lim_{x \rightarrow a} [c \cdot f(x)] = c \cdot \lim_{x \rightarrow a} f(x)$
- $\lim_{x \rightarrow a} [f(x) \cdot g(x)] = \lim_{x \rightarrow a} f(x) \cdot \lim_{x \rightarrow a} g(x)$
- $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)}$ if $\lim_{x \rightarrow a} g(x) \neq 0$




Use these graphs to estimate:

- $$\begin{aligned} 1. \lim_{x \rightarrow -2} [f(x) + 5g(x)] &= \lim_{x \rightarrow -2} f(x) + \lim_{x \rightarrow -2} [5g(x)] \\ &= \lim_{x \rightarrow -2} f(x) + 5 \lim_{x \rightarrow -2} g(x) \\ &= 2 + 5(-1) \\ &= -3 \end{aligned}$$

Calculating Limits using Limit Laws

1. $\lim_{x \rightarrow a} [f(x) + g(x)] = \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x)$
2. $\lim_{x \rightarrow a} [f(x) - g(x)] = \lim_{x \rightarrow a} f(x) - \lim_{x \rightarrow a} g(x)$
3. $\lim_{x \rightarrow a} [c \cdot f(x)] = c \cdot \lim_{x \rightarrow a} f(x)$
4. $\lim_{x \rightarrow a} [f(x) \cdot g(x)] = \lim_{x \rightarrow a} f(x) \cdot \lim_{x \rightarrow a} g(x)$
5. $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)}$ if $\lim_{x \rightarrow a} g(x) \neq 0$




Use these graphs to estimate:

2. $\lim_{x \rightarrow 1} [f(x)g(x)]$
 $= \lim_{x \rightarrow 1} f(x) \cdot \lim_{x \rightarrow 1} g(x)$
 $\leftarrow \lim_{x \rightarrow 1} g(x)$ does not exist
(we cannot use the limit laws)

Calculating Limits using Limit Laws

- $\lim_{x \rightarrow a} [f(x) + g(x)] = \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x)$
- $\lim_{x \rightarrow a} [f(x) - g(x)] = \lim_{x \rightarrow a} f(x) - \lim_{x \rightarrow a} g(x)$
- $\lim_{x \rightarrow a} [c \cdot f(x)] = c \cdot \lim_{x \rightarrow a} f(x)$
- $\lim_{x \rightarrow a} [f(x) \cdot g(x)] = \lim_{x \rightarrow a} f(x) \cdot \lim_{x \rightarrow a} g(x)$
- $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)}$ if $\lim_{x \rightarrow a} g(x) \neq 0$



Use these graphs to estimate:


$$\begin{aligned} 2a. \quad \lim_{x \rightarrow 1^-} [f(x)g(x)] &= \lim_{x \rightarrow 1^-} f(x) \cdot \lim_{x \rightarrow 1^-} g(x) \\ &= 2 \cdot -2 = -4 \end{aligned}$$

$$\begin{aligned} 2b. \quad \lim_{x \rightarrow 1^+} [f(x)g(x)] &= \lim_{x \rightarrow 1^+} f(x) \cdot \lim_{x \rightarrow 1^+} g(x) \\ &= 2 \cdot -1 = -2 \end{aligned}$$

$\Rightarrow \lim_{x \rightarrow 1} [f(x)g(x)]$ does not exist

Calculating Limits using Limit Laws

1. $\lim_{x \rightarrow a} [f(x) + g(x)] = \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x)$
2. $\lim_{x \rightarrow a} [f(x) - g(x)] = \lim_{x \rightarrow a} f(x) - \lim_{x \rightarrow a} g(x)$
3. $\lim_{x \rightarrow a} [c \cdot f(x)] = c \cdot \lim_{x \rightarrow a} f(x)$
4. $\lim_{x \rightarrow a} [f(x) \cdot g(x)] = \lim_{x \rightarrow a} f(x) \cdot \lim_{x \rightarrow a} g(x)$
5. $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)}$ if $\lim_{x \rightarrow a} g(x) \neq 0$



Use these graphs to estimate:


$$3. \lim_{x \rightarrow 2} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow 2} f(x)}{\lim_{x \rightarrow 2} g(x)}$$

⚡ $\lim_{x \rightarrow 2} g(x) = 0$

(we cannot use the limit laws)

Calculating Limits using Limit Laws

1. $\lim_{x \rightarrow a} [f(x) + g(x)] = \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x)$
2. $\lim_{x \rightarrow a} [f(x) - g(x)] = \lim_{x \rightarrow a} f(x) - \lim_{x \rightarrow a} g(x)$
3. $\lim_{x \rightarrow a} [c \cdot f(x)] = c \cdot \lim_{x \rightarrow a} f(x)$
4. $\lim_{x \rightarrow a} [f(x) \cdot g(x)] = \lim_{x \rightarrow a} f(x) \cdot \lim_{x \rightarrow a} g(x)$
5. $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)}$ if $\lim_{x \rightarrow a} g(x) \neq 0$



Use these graphs to estimate:

Lets try without limit laws:

- 3a. $\lim_{x \rightarrow 2^-} \frac{f(x)}{g(x)} = -\infty$
since $\lim_{x \rightarrow 2^-} f(x) \approx 1.6$, and $g(x)$ approaches 0, $g(x) < 0$
- 3b. $\lim_{x \rightarrow 2^+} \frac{f(x)}{g(x)} = \infty$
since $\lim_{x \rightarrow 2^+} f(x) \approx 1.6$, and $g(x)$ approaches 0, $g(x) > 0$

More Limits Laws

1. $\lim_{x \rightarrow a} [f(x) + g(x)] = \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x)$
2. $\lim_{x \rightarrow a} [f(x) - g(x)] = \lim_{x \rightarrow a} f(x) - \lim_{x \rightarrow a} g(x)$
3. $\lim_{x \rightarrow a} [c \cdot f(x)] = c \cdot \lim_{x \rightarrow a} f(x)$
4. $\lim_{x \rightarrow a} [f(x) \cdot g(x)] = \lim_{x \rightarrow a} f(x) \cdot \lim_{x \rightarrow a} g(x)$
5. $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)}$ if $\lim_{x \rightarrow a} g(x) \neq 0$
6. $\lim_{x \rightarrow a} [f(x)]^n = [\lim_{x \rightarrow a} f(x)]^n$ for n a positive integer
7. $\lim_{x \rightarrow a} c = c$
8. $\lim_{x \rightarrow a} x^n = a^n$
9. $\lim_{x \rightarrow a} \sqrt[n]{x} = \sqrt[n]{a}$ for n a positive integer
(if n is even we require $a > 0$)
10. $\lim_{x \rightarrow a} \sqrt[n]{f(x)} = \sqrt[n]{\lim_{x \rightarrow a} f(x)}$ for n a positive integer
(if n is even we require $\lim_{x \rightarrow a} f(x) > 0$)

Limit Laws: Examples

1. $\lim_{x \rightarrow a} [f(x) + g(x)] = \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x)$
2. $\lim_{x \rightarrow a} [f(x) - g(x)] = \lim_{x \rightarrow a} f(x) - \lim_{x \rightarrow a} g(x)$
3. $\lim_{x \rightarrow a} [c \cdot f(x)] = c \cdot \lim_{x \rightarrow a} f(x)$
4. $\lim_{x \rightarrow a} [f(x) \cdot g(x)] = \lim_{x \rightarrow a} f(x) \cdot \lim_{x \rightarrow a} g(x)$
5. $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)}$ if $\lim_{x \rightarrow a} g(x) \neq 0$
6. $\lim_{x \rightarrow a} [f(x)]^n = [\lim_{x \rightarrow a} f(x)]^n$ for n a positive integer
7. $\lim_{x \rightarrow a} c = c$
8. $\lim_{x \rightarrow a} x^n = a^n$
9. $\lim_{x \rightarrow a} \sqrt[n]{x} = \sqrt[n]{a}$ for n a positive integer
(if n is even we require $a > 0$)
10. $\lim_{x \rightarrow a} \sqrt[n]{f(x)} = \sqrt[n]{\lim_{x \rightarrow a} f(x)}$ for n a positive integer
(if n is even we require $\lim_{x \rightarrow a} f(x) > 0$)

$$\lim_{x \rightarrow 5} (2x^2 - 3x + 4) = \lim_{x \rightarrow 5} (2x^2) - \lim_{x \rightarrow 5} (3x) + \lim_{x \rightarrow 5} 4 \quad (\text{law 1 and 2})$$

$$= 2 \lim_{x \rightarrow 5} (x^2) - 3 \lim_{x \rightarrow 5} (x) + 4 \quad (\text{law 3 and 7})$$

$$= 2 \cdot 5^2 - 3 \cdot 5 + 4 = 39 \quad (\text{law 8})$$

Limit Laws: Examples

$$\begin{aligned}\lim_{x \rightarrow -2} \frac{x^3 + 2x^2 - 1}{5 - 3x} &= \frac{\lim_{x \rightarrow -2} (x^3 + 2x^2 - 1)}{\lim_{x \rightarrow -2} (5 - 3x)} && \text{(law 5)} \\ &= \frac{\lim_{x \rightarrow -2} x^3 + 2 \lim_{x \rightarrow -2} x^2 - \lim_{x \rightarrow -2} 1}{\lim_{x \rightarrow -2} 5 - 3 \lim_{x \rightarrow -2} x} && \text{(law 1, 2, 3)} \\ &= \frac{(-2)^3 + 2 \cdot (-2)^2 - 1}{5 - 3 \cdot (-2)} && \text{(law 7, 8)} \\ &= -\frac{1}{11}\end{aligned}$$

Computing Limits: Direct Substitution Property

Direct Substitution Property

If f is a polynomial or a rational and a is in the domain of f , then:

$$\lim_{x \rightarrow a} f(x) = f(a)$$

Works also for one-sided limits $\lim_{x \rightarrow a^\pm} f(x) = f(a)$.

Works also for algebraic functions if $f(x)$ is defined close to a .

The function $f(x) = 2x^2 - 3x + 4$ is a polynomial and hence:

$$\lim_{x \rightarrow 5} f(x) = f(5) = 2 \cdot 5^2 - 3 \cdot 5 + 4 = 39$$

The function $g(x) = \frac{x^3 + 2x^2 - 1}{5 - 3x}$ is rational and -2 is in the domain; hence:

$$\lim_{x \rightarrow -2} g(x) = g(-2) = \frac{(-2)^3 + 2 \cdot (-2)^2 - 1}{5 - 3 \cdot (-2)} = -\frac{1}{11}$$

Computing Limits: Function Replacement

Function Replacement

If $f(x) = g(x)$ for all $x \neq a$, then $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} g(x)$
(provided that the limit exists).

Actually it suffices $f(x) = g(x)$ when x is close to a .

Find $\lim_{x \rightarrow 1} \frac{x^2 - 1}{x - 1}$.

- ▶ Direct substitution is not applicable because $x = 1$ is not in the domain.

We replace the function:

$$\frac{x^2 - 1}{x - 1} = \frac{(x + 1)(x - 1)}{x - 1} \text{ for } x \neq 1 = x + 1$$

As a consequence

$$\lim_{x \rightarrow 1} \frac{x^2 - 1}{x - 1} = \lim_{x \rightarrow 1} x + 1 = 1 + 1 = 2$$

Computing Limits: Function Replacement

Find $\lim_{x \rightarrow 1} g(x)$ where

$$g(x) = \begin{cases} 2x + 1 & \text{for } x \neq 1, \\ \pi & \text{for } x = 1 \end{cases}$$

We have:

$$g(x) = 2x + 1 \quad \text{for all } x \neq 1$$

As a consequence:

$$\lim_{x \rightarrow 1} g(x) = \lim_{x \rightarrow 1} 2x + 1 = 2 + 1 = 3$$

Computing Limits: Function Replacement

Find

$$\lim_{h \rightarrow 0} \frac{(3+h)^2 - 9}{h}$$

We have:

$$\frac{(3+h)^2 - 9}{h} = \frac{9 + 6h + h^2 - 9}{h} = \frac{6h + h^2}{h} \stackrel{\text{for } h \neq 0}{=} 6 + h$$

As a consequence:

$$\lim_{h \rightarrow 0} \frac{(3+h)^2 - 9}{h} = \lim_{h \rightarrow 0} (6 + h) = 6$$

Computing Limits: Function Replacement

Find

$$\lim_{t \rightarrow 0} \frac{\sqrt{t^2 + 9} - 3}{t^2}$$

We have:

$$\begin{aligned} \frac{\sqrt{t^2 + 9} - 3}{t^2} &= \frac{\sqrt{t^2 + 9} - 3}{t^2} \cdot \frac{\sqrt{t^2 + 9} + 3}{\sqrt{t^2 + 9} + 3} = \frac{t^2 + 9 - 9}{t^2 \cdot (\sqrt{t^2 + 9} + 3)} \\ &= \frac{t^2}{t^2 \cdot (\sqrt{t^2 + 9} + 3)} \stackrel{\text{for } t \neq 0}{=} \frac{1}{\sqrt{t^2 + 9} + 3} \end{aligned}$$

As a consequence:

$$\begin{aligned} \lim_{t \rightarrow 0} \frac{\sqrt{t^2 + 9} - 3}{t^2} &= \lim_{t \rightarrow 0} \frac{1}{\sqrt{t^2 + 9} + 3} \\ &= \frac{1}{\sqrt{\lim_{t \rightarrow 0} (t^2 + 9)} + 3} \quad \text{by laws 5, 1, 9, 7} \\ &= \frac{1}{\sqrt{9 + 3}} = \frac{1}{6} \end{aligned}$$

Limits and One-Sided Limits

We recall the following theorem:

$$\lim_{x \rightarrow a} f(x) = L \quad \text{if and only if} \quad \lim_{x \rightarrow a^-} f(x) = L = \lim_{x \rightarrow a^+} f(x)$$

The theorem in words:

- ▶ The limit of $f(x)$, for x approaching a , is L if and only if the left-limit and the right-limit at a are both L .

The limit laws also apply for one-sided limits!

- ▶ if $\lim_{x \rightarrow a^-} f(x) \neq \lim_{x \rightarrow a^+} f(x)$
then $\lim_{x \rightarrow a} f(x)$ does not exist

Computing Limits: Function Replacement

Function replacement for one-sided limits:

If $f(x) = g(x)$ for all $x < a$, then $\lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^-} g(x)$.

If $f(x) = g(x)$ for all $x > a$, then $\lim_{x \rightarrow a^+} f(x) = \lim_{x \rightarrow a^+} g(x)$.

Find $\lim_{x \rightarrow 2^-} g(x)$ where

$$g(x) = \begin{cases} x^2 & \text{for } x < 2 \\ 5x + 1 & \text{for } x \geq 2 \end{cases}$$

We have

$$g(x) = x^2 \quad \text{for all } x < 2$$

Hence:

$$\lim_{x \rightarrow 2^-} g(x) = \lim_{x \rightarrow 2^-} x^2 = 4$$

Computing Limits: Function Replacement

For one-sided limits we have:

If $f(x) = g(x)$ for all $x < a$, then $\lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^-} g(x)$.

If $f(x) = g(x)$ for all $x > a$, then $\lim_{x \rightarrow a^+} f(x) = \lim_{x \rightarrow a^+} g(x)$.

Find $\lim_{x \rightarrow 0} |x|$ where

$$|x| = \begin{cases} x & \text{for } x \geq 0 \\ -x & \text{for } x < 0 \end{cases}$$

Since $|x| = x$ for all $x > 0$ we obtain:

$$\lim_{x \rightarrow 0^+} |x| = \lim_{x \rightarrow 0^+} x = 0$$

Since $|x| = -x$ for all $x < 0$ we obtain:

$$\lim_{x \rightarrow 0^-} |x| = \lim_{x \rightarrow 0^-} -x = 0$$

Hence $\lim_{x \rightarrow 0} |x| = 0$.

Computing Limits: Function Replacement

For one-sided limits we have:

If $f(x) = g(x)$ for all $x < a$, then $\lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^-} g(x)$.

If $f(x) = g(x)$ for all $x > a$, then $\lim_{x \rightarrow a^+} f(x) = \lim_{x \rightarrow a^+} g(x)$.

Proof that $\lim_{x \rightarrow 0} \frac{|x|}{x}$ does not exist.

For all $x > 0$ we have $\frac{|x|}{x} = \frac{x}{x} = 1$. Thus

$$\lim_{x \rightarrow 0^+} \frac{|x|}{x} = \lim_{x \rightarrow 0^+} 1 = 1$$

For all $x < 0$ we have $\frac{|x|}{x} = \frac{-x}{x} = -1$. Thus

$$\lim_{x \rightarrow 0^-} \frac{|x|}{x} = \lim_{x \rightarrow 0^-} -1 = -1$$

Hence $\lim_{x \rightarrow 0} \frac{|x|}{x}$ does not exist since $\lim_{x \rightarrow 0^-} \frac{|x|}{x} \neq \lim_{x \rightarrow 0^+} \frac{|x|}{x}$.

Properties of Limits

If

- ▶ $f(x) \leq g(x)$ when x is near a (except possibly a),
- ▶ $\lim_{x \rightarrow a} f(x)$ exists, and
- ▶ $\lim_{x \rightarrow a} g(x)$ exist,

then

$$\lim_{x \rightarrow a} f(x) \leq \lim_{x \rightarrow a} g(x)$$

Formally, near a means on $(a - \epsilon, a + \epsilon) \setminus \{a\}$ for some $\epsilon > 0$.

We have $x^3 \leq x^2$ for $x \in (-1, 1)$.

As a consequence:

$$\lim_{x \rightarrow a} x^3 \leq \lim_{x \rightarrow a} x^2$$

for all $a \in (-1, 1)$.

Properties of Limits


The Squeeze Theorem

If $f(x) \leq g(x) \leq h(x)$ when x is near a (except possibly a) and

$$\lim_{x \rightarrow a} f(x) = L = \lim_{x \rightarrow a} h(x)$$

then

$$\lim_{x \rightarrow a} g(x) = L$$



Here f is below g , and h is above g (close to a). If f and h have the same limit, then the squeezed function g also has.

Properties of Limits

Show that $\lim_{x \rightarrow 0} g(x) = 0$ where $g(x) = x^2 \cdot \sin \frac{1}{x}$.

The application of limit laws

$$\lim_{x \rightarrow 0} (x^2 \cdot \sin \frac{1}{x}) = (\lim_{x \rightarrow 0} x^2) \cdot (\lim_{x \rightarrow 0} \sin \frac{1}{x})$$

does not work since $\lim_{x \rightarrow 0} \sin \frac{1}{x}$ does not exist.

To apply the squeeze theorem we need:

- ▶ a function f smaller (\leq) than g , and
- ▶ a function h bigger (\geq) than g

for which $\lim_{x \rightarrow 0} f(x) = 0$ and $\lim_{x \rightarrow 0} h(x) = 0$.

We know that $-1 \leq \sin \frac{1}{x} \leq 1$ and hence


$$-x^2 \leq x^2 \cdot \sin \frac{1}{x} \leq x^2$$

Properties of Limits

We have

$$-x^2 \leq x^2 \cdot \sin \frac{1}{x} \leq x^2$$

We take $f(x) = -x^2$ and $h(x) = x^2$.



We know $\lim_{x \rightarrow 0} x^2 = 0$ and $\lim_{x \rightarrow 0} -x^2 = 0$.

Hence by the squeeze theorem we get: $\lim_{x \rightarrow 0} x^2 \cdot \sin \frac{1}{x} = 0$.