



- Lecture 1: Introduction, Abstract Rewriting
- Lecture 2: Term Rewriting
- Lecture 3: Combinatory Logic
- Lecture 4: Termination
- Lecture 5: Matching, Unification
- Lecture 6: Equational Reasoning, Completion
- Lecture 7: Confluence
- Lecture 8: Modularity
- Lecture 9: Strategies
- Lecture 10: Decidability
- Lecture 11: **Infinitary Rewriting**

Outline



- Overview
- Infinitary Rewriting

Infinitary Rewriting

Example (The Stream of Natural Numbers)



Infinite Terms





Definition

An **infinite term** is a partial map $t : \mathbb{N}^* \rightarrow \Sigma$ from positions to symbols such that:

- $t(\epsilon) \in \Sigma$, and
- $t(ip) \in \Sigma \iff 1 \leq i \leq \#(t(p))$

Infinite Terms



Definition

An **infinite term** is a partial map $t : \mathbb{N}^* \rightharpoonup \Sigma$ from positions to symbols such that:

- $t(\epsilon) \in \Sigma$, and
- $t(ip) \in \Sigma \iff 1 \leq i \leq \#(t(p))$

The **set of finite and infinite terms** is denoted by $\mathcal{T}^\infty(\Sigma, \mathcal{X})$.

Infinite Terms as Metric Space

Definition

We define a **metric** d on $\mathcal{T}^\infty(\Sigma, \mathcal{X})$ by:

$$d(s, t) = 2^{-|p|} \text{ where } p \text{ is the highest position such that } s(p) \neq t(p)$$

The first difference is at depth 2, hence $d(s, t) = 2^{-2} = 0.25$.

Infinite Terms as Metric Space

Definition

We define a **metric** d on $\mathcal{T}^\infty(\Sigma, \mathcal{X})$ by:

$$d(s, t) = 2^{-|p|} \text{ where } p \text{ is the highest position such that } s(p) \neq t(p)$$

Note that $d(s, t) = 0 \iff s = t$.

The first difference is at depth 2, hence $d(s, t) = 2^{-2} = 0.25$.

Infinite Terms as Metric Space


Definition

We define a **metric d** on $\mathcal{T}^\infty(\Sigma, \mathcal{X})$ by:

$$d(s, t) = 2^{-|p|} \text{ where } p \text{ is the highest position such that } s(p) \neq t(p)$$

Note that $d(s, t) = 0 \iff s = t$.

Example



Infinite Terms as Metric Space


Definition

We define a **metric d** on $\mathcal{T}^\infty(\Sigma, \mathcal{X})$ by:

$$d(s, t) = 2^{-|p|} \text{ where } p \text{ is the highest position such that } s(p) \neq t(p)$$

Note that $d(s, t) = 0 \iff s = t$.

Example



The first difference is at depth 2, hence $d(s, t) = 2^{-2} = 0.25$.

Infinitary Rewriting

Example

$$f(x, x) \rightarrow f(a, b)$$

$$a \rightarrow c(a)$$

$$b \rightarrow c(b)$$

Infinitary Rewriting

Example

$$f(x, x) \rightarrow f(a, b)$$

$$a \rightarrow c(a)$$

$$b \rightarrow c(b)$$

$$f(a, b)$$

Infinitary Rewriting

Example

$$f(x, x) \rightarrow f(a, b)$$

$$a \rightarrow c(a)$$

$$b \rightarrow c(b)$$

$$f(a, b) \rightarrow f(c(a), b)$$

Infinitary Rewriting

Example

$$f(x, x) \rightarrow f(a, b)$$

$$a \rightarrow c(a)$$

$$b \rightarrow c(b)$$

$$\begin{aligned} f(a, b) &\rightarrow f(c(a), b) \\ &\rightarrow f(c(c(a)), b) \end{aligned}$$

Infinitary Rewriting

Example

$$f(x, x) \rightarrow f(a, b)$$

$$a \rightarrow c(a)$$

$$b \rightarrow c(b)$$

$$f(a, b) \rightarrow f(c(a), b)$$

$$\rightarrow f(c(c(a)), b)$$

$$\rightarrow\!\!\!\rightarrow f(c(c(c(\dots))), b)$$

Infinitary Rewriting

Example

$$f(x, x) \rightarrow f(a, b)$$

$$a \rightarrow c(a)$$

$$b \rightarrow c(b)$$

$$f(a, b) \rightarrow f(c(a), b)$$

$$\rightarrow f(c(c(a)), b)$$

$$\rightarrow\!\!\!\rightarrow f(c(c(c(\dots))), b)$$

$$\rightarrow f(c(c(c(\dots))), c(b))$$

Infinitary Rewriting

Example

$$f(x, x) \rightarrow f(a, b)$$

$$a \rightarrow c(a)$$

$$b \rightarrow c(b)$$

$$f(a, b) \rightarrow f(c(a), b)$$

$$\rightarrow f(c(c(a)), b)$$

$$\rightarrow\!\!\!\rightarrow f(c(c(c(\dots))), b)$$

$$\rightarrow f(c(c(c(\dots))), c(b))$$

$$\rightarrow f(c(c(c(\dots))), c(c(b)))$$

Infinitary Rewriting

Example

$$f(x, x) \rightarrow f(a, b)$$

$$a \rightarrow c(a)$$

$$b \rightarrow c(b)$$

$$f(a, b) \rightarrow f(c(a), b)$$

$$\rightarrow f(c(c(a)), b)$$

$$\rightarrow\!\!\!\rightarrow f(c(c(c(\dots))), b)$$

$$\rightarrow f(c(c(c(\dots))), c(b))$$

$$\rightarrow f(c(c(c(\dots))), c(c(b)))$$

$$\rightarrow\!\!\!\rightarrow f(c(c(c(\dots))), c(c(c(\dots))))$$

Infinitary Rewriting

Example

$$f(x, x) \rightarrow f(a, b)$$

$$a \rightarrow c(a)$$

$$b \rightarrow c(b)$$

$$f(a, b) \rightarrow f(c(a), b)$$

$$\rightarrow f(c(c(a)), b)$$

$$\rightarrow\!\!\!\rightarrow f(c(c(c(\dots))), b)$$

$$\rightarrow f(c(c(c(\dots))), c(b))$$

$$\rightarrow f(c(c(c(\dots))), c(c(b)))$$

$$\rightarrow\!\!\!\rightarrow f(c(c(c(\dots))), c(c(c(\dots))))$$

$$\rightarrow f(a, b)$$

Infinitary Rewriting

Example

$$f(x, x) \rightarrow f(a, b)$$

$$a \rightarrow c(a)$$

$$b \rightarrow c(b)$$

$$f(a, b) \rightarrow f(c(a), b)$$

$$\rightarrow f(c(c(a)), b)$$

$$\rightarrow\!\!\!\rightarrow f(c(c(c(\dots))), b)$$

$$\rightarrow f(c(c(c(\dots))), c(b))$$

$$\rightarrow f(c(c(c(\dots))), c(c(b)))$$

$$\rightarrow\!\!\!\rightarrow f(c(c(c(\dots))), c(c(c(\dots))))$$

$$\rightarrow f(a, b)$$

$$\rightarrow \dots$$

Infinitary Rewriting

Example

$$f(x, x) \rightarrow f(a, b)$$

$$a \rightarrow c(a)$$

$$b \rightarrow c(b)$$

$$f(a, b) \rightarrow f(c(a), b)$$

$$\rightarrow f(c(c(a)), b)$$

$$\rightarrow\!\!\!\rightarrow f(c(c(c(\dots))), b)$$

$$\rightarrow f(c(c(c(\dots))), c(b))$$

$$\rightarrow f(c(c(c(\dots))), c(c(b)))$$

$$\rightarrow\!\!\!\rightarrow f(c(c(c(\dots))), c(c(c(\dots))))$$


$$\rightarrow f(a, b)$$

$$\rightarrow \dots$$

We need transfinite reductions...

Ordinals

$0, 1, 2, \dots, \omega, \omega + 1, \omega + 2, \dots, \omega + \omega = \omega \cdot 2, \dots, \omega \cdot 3, \dots, \omega^2, \dots, \omega^\omega, \dots$



Note that ω is the smallest infinite ordinal.

Ordinals

Definition

A set S is **transitive** if $x \in S$ implies $x \subseteq S$.

Ordinals

Definition

A set S is **transitive** if $x \in S$ implies $x \subseteq S$.

An **ordinal** is a transitive set whose elements are transitive sets.

Ordinals

Definition

A set S is **transitive** if $x \in S$ implies $x \subseteq S$.

An **ordinal** is a transitive set whose elements are transitive sets.

Example $(0, 1, 2, 3, \dots)$

$\emptyset, \{\emptyset\}, \{\emptyset, \{\emptyset\}\}, \{\emptyset, \{\emptyset\}, \{\emptyset, \{\emptyset\}\}\}, \dots$

Ordinals

Definition

A set S is **transitive** if $x \in S$ implies $x \subseteq S$.

An **ordinal** is a transitive set whose elements are transitive sets.

Example $(0, 1, 2, 3, \dots)$

$\emptyset, \{\emptyset\}, \{\emptyset, \{\emptyset\}\}, \{\emptyset, \{\emptyset\}, \{\emptyset, \{\emptyset\}\}\}, \dots$

Definition

We define $\alpha < \beta \iff \alpha \in \beta$.

Ordinals

Definition

A set S is **transitive** if $x \in S$ implies $x \subseteq S$.

An **ordinal** is a transitive set whose elements are transitive sets.

Example $(0, 1, 2, 3, \dots)$

$\emptyset, \{\emptyset\}, \{\emptyset, \{\emptyset\}\}, \{\emptyset, \{\emptyset\}, \{\emptyset, \{\emptyset\}\}\}, \dots$

Definition

We define $\alpha < \beta \iff \alpha \in \beta$.

Lemma

The relation $<$ is a total order on ordinals.

Ordinals

Definition

A set S is **transitive** if $x \in S$ implies $x \subseteq S$.

An **ordinal** is a transitive set whose elements are transitive sets.

Example $(0, 1, 2, 3, \dots)$

$\emptyset, \{\emptyset\}, \{\emptyset, \{\emptyset\}\}, \{\emptyset, \{\emptyset\}, \{\emptyset, \{\emptyset\}\}\}, \dots$

Definition

We define $\alpha < \beta \iff \alpha \in \beta$.

Lemma

The relation $<$ is a total order on ordinals.


Lemma

For every ordinal β , we have $\beta = \{\alpha \mid \alpha < \beta\}$.

Ordinals

Definition

For ordinals α , we define $\alpha^+ = \alpha \cup \{\alpha\}$, the **successor of α** .




Ordinals

Definition

For ordinals α , we define $\alpha^+ = \alpha \cup \{\alpha\}$, the **successor of α** .

An ordinal α is a **successor** ordinal if $\alpha = \beta^+$ for some ordinal β .



Example

Successor ordinals: $1, 2, \omega + 1, \omega \cdot 3 + 2, \dots$


Ordinals

Definition

For ordinals α , we define $\alpha^+ = \alpha \cup \{\alpha\}$, the **successor of α** .

An ordinal α is a **successor** ordinal if $\alpha = \beta^+$ for some ordinal β .

If $\alpha \neq 0$ and α is not a successor ordinal, then α is called **limit** ordinal.



Example

Successor ordinals: $1, 2, \omega + 1, \omega \cdot 3 + 2, \dots$ Limit ordinals: $\omega, \omega \cdot 2, \omega \cdot 3, \omega^2, \dots$

Transfinite Reductions

Example

$$f(x, x) \rightarrow f(a, b)$$

$$a \rightarrow c(a)$$

$$b \rightarrow c(b)$$

Transfinite Reductions

Example

$$f(x, x) \rightarrow f(a, b)$$

$$a \rightarrow c(a)$$

$$b \rightarrow c(b)$$

A reduction of length $\omega \cdot 2 + 1$:

$$f(a, b)$$

Transfinite Reductions

Example

$$f(x, x) \rightarrow f(a, b)$$

$$a \rightarrow c(a)$$

$$b \rightarrow c(b)$$

A reduction of length $\omega \cdot 2 + 1$:

$$f(a, b) \rightarrow^\omega f(c^\omega, b)$$

Transfinite Reductions

Example

$$f(x, x) \rightarrow f(a, b)$$

$$a \rightarrow c(a)$$

$$b \rightarrow c(b)$$

A reduction of length $\omega \cdot 2 + 1$:

$$f(a, b) \rightarrow^\omega f(c^\omega, b) \rightarrow^\omega f(c^\omega, c^\omega)$$

Transfinite Reductions

Example

$$f(x, x) \rightarrow f(a, b)$$

$$a \rightarrow c(a)$$

$$b \rightarrow c(b)$$

A reduction of length $\omega \cdot 2 + 1$:

$$f(a, b) \rightarrow^\omega f(c^\omega, b) \rightarrow^\omega f(c^\omega, c^\omega) \rightarrow f(a, b)$$

Transfinite Reductions

Example

$$f(x, x) \rightarrow f(a, b)$$

$$a \rightarrow c(a)$$

$$b \rightarrow c(b)$$

A reduction of length $\omega \cdot 2 + 1$:

$$f(a, b) \rightarrow^\omega f(c^\omega, b) \rightarrow^\omega f(c^\omega, c^\omega) \rightarrow f(a, b)$$

A reduction of length $\omega + 1$:

$$f(a, b)$$

Transfinite Reductions

Example

$$f(x, x) \rightarrow f(a, b)$$

$$a \rightarrow c(a)$$

$$b \rightarrow c(b)$$

A reduction of length $\omega \cdot 2 + 1$:

$$f(a, b) \rightarrow^\omega f(c^\omega, b) \rightarrow^\omega f(c^\omega, c^\omega) \rightarrow f(a, b)$$

A reduction of length $\omega + 1$:

$$f(a, b) \rightarrow^\omega f(c^\omega, c^\omega)$$

Transfinite Reductions

Example

$$f(x, x) \rightarrow f(a, b)$$

$$a \rightarrow c(a)$$

$$b \rightarrow c(b)$$

A reduction of length $\omega \cdot 2 + 1$:

$$f(a, b) \rightarrow^\omega f(c^\omega, b) \rightarrow^\omega f(c^\omega, c^\omega) \rightarrow f(a, b)$$

A reduction of length $\omega + 1$:

$$f(a, b) \rightarrow^\omega f(c^\omega, c^\omega) \rightarrow f(a, b)$$

Transfinite Reductions

Example

$$f(x, x) \rightarrow f(a, b)$$

$$a \rightarrow c(a)$$

$$b \rightarrow c(b)$$

A reduction of length $\omega \cdot 2 + 1$:

$$f(a, b) \rightarrow^\omega f(c^\omega, b) \rightarrow^\omega f(c^\omega, c^\omega) \rightarrow f(a, b)$$

A reduction of length $\omega + 1$:

$$f(a, b) \rightarrow^\omega f(c^\omega, c^\omega) \rightarrow f(a, b)$$

by alternating $f(a, b) \rightarrow f(c(a), b) \rightarrow f(c(a), c(b)) \rightarrow \dots$

Transfinite Reductions

Definition

Let α be an ordinal, and $\tau : (t_\beta \rightarrow t_{\beta+1})_{\beta < \alpha}$ a sequence of reduction steps.

Transfinite Reductions

Definition

Let α be an ordinal, and $\tau : (t_\beta \rightarrow t_{\beta+1})_{\beta < \alpha}$ a sequence of reduction steps.

We use d_β to denote the depth of the rewrite step $t_\beta \rightarrow t_{\beta+1}$.

Transfinite Reductions

Definition

Let α be an ordinal, and $\tau : (t_\beta \rightarrow t_{\beta+1})_{\beta < \alpha}$ a sequence of reduction steps.

We use d_β to denote the depth of the rewrite step $t_\beta \rightarrow t_{\beta+1}$.

Then τ is an **infinite reduction** of length α if for every limit ordinal $\lambda < \alpha$:

- 1 the distance $d(t_\beta, t_\lambda)$ tends to 0, $\forall \epsilon > 0. \exists \beta < \lambda. \forall \beta < \gamma < \lambda. d(t_\gamma, t_\lambda) \leq \epsilon$
- 2 the depth d_β tends to infinity $\forall n. \exists \beta < \lambda. \forall \beta < \gamma < \lambda. d_\gamma \geq n$

as β approaches λ from below.

Transfinite Reductions

Definition

Let α be an ordinal, and $\tau : (t_\beta \rightarrow t_{\beta+1})_{\beta < \alpha}$ a sequence of reduction steps.

We use d_β to denote the depth of the rewrite step $t_\beta \rightarrow t_{\beta+1}$.

Then τ is an **infinite reduction** of length α if for every limit ordinal $\lambda < \alpha$:

- 1 the distance $d(t_\beta, t_\lambda)$ tends to 0, $\forall \epsilon > 0. \exists \beta < \lambda. \forall \beta < \gamma < \lambda. d(t_\gamma, t_\lambda) \leq \epsilon$
- 2 the depth d_β tends to infinity $\forall n. \exists \beta < \lambda. \forall \beta < \gamma < \lambda. d_\gamma \geq n$

as β approaches λ from below.

Example

Let $R = \{a \rightarrow a, b \rightarrow b\}$. Condition (1) excludes **jumps in the limit**:

$$a \rightarrow a \rightarrow a \rightarrow \dots \underbrace{b}_{t_\omega} \rightarrow b \rightarrow \dots$$

Transfinite Reductions

Definition

Let α be an ordinal, and $\tau : (t_\beta \rightarrow t_{\beta+1})_{\beta < \alpha}$ a sequence of reduction steps.

We use d_β to denote the depth of the rewrite step $t_\beta \rightarrow t_{\beta+1}$.

Then τ is an **infinite reduction** of length α if for every limit ordinal $\lambda < \alpha$:

- 1 the distance $d(t_\beta, t_\lambda)$ tends to 0, $\forall \epsilon > 0. \exists \beta < \lambda. \forall \beta < \gamma < \lambda. d(t_\gamma, t_\lambda) \leq \epsilon$
- 2 the depth d_β tends to infinity $\forall n. \exists \beta < \lambda. \forall \beta < \gamma < \lambda. d_\gamma \geq n$

as β approaches λ from below.


Example (We want more than Cauchy-convergence...)

Let $R = \{f(x) \rightarrow f(c(x))\}$. Condition (2) excludes sequences like:

$$f(a) \rightarrow_\epsilon f(c(a)) \rightarrow_\epsilon f(c(c(a))) \rightarrow_\epsilon \dots \rightarrow_\epsilon^\omega f(c^\omega) \rightarrow \dots$$

where the **activity does not move downwards**.

Transfinite Reductions Visualized



The blue lines indicate the depth of the activity/rewrite steps.

The activity tends to infinity when approaching limit ordinals.

Why more than Cauchy-convergence?

We consider the TRS:

$$f(x, y) \rightarrow f(y, x)$$

$$a \rightarrow b$$

Why more than Cauchy-convergence?

We consider the TRS:

$$\begin{aligned} f(x, y) &\rightarrow f(y, x) \\ a &\rightarrow b \end{aligned}$$

We start from $f(a, a)$ and trace the left occurrence of a :

Why more than Cauchy-convergence?

We consider the TRS:

$$\begin{aligned} f(x, y) &\rightarrow f(y, x) \\ a &\rightarrow b \end{aligned}$$

We start from $f(a, a)$ and trace the left occurrence of a :

$$f(\bar{a}, a) \rightarrow f(a, \bar{a}) \rightarrow f(\bar{a}, a) \rightarrow f(a, \bar{a}) \rightarrow^\omega ?$$

The rewrite sequence without overlining is Cauchy-convergent.

However, what are the residuals of the left a after ω -many steps?

Why more than Cauchy-convergence?

We consider the TRS:

$$\begin{aligned} f(x, y) &\rightarrow f(y, x) \\ a &\rightarrow b \end{aligned}$$

We start from $f(a, a)$ and trace the left occurrence of a :

$$f(\bar{a}, a) \rightarrow f(a, \bar{a}) \rightarrow f(\bar{a}, a) \rightarrow f(a, \bar{a}) \rightarrow^\omega ?$$

The rewrite sequence without overlining is Cauchy-convergent.

However, what are the residuals of the left a after ω -many steps?

Although it appears as if the term has a limit, this is only a syntactic accident. The subterms get swapped all the time. . .

Definition

A reduction of length α is **strongly convergent** if for every limit ordinal $\lambda \leq \alpha$ the depth d_β tends to infinity as β approaches λ from below, and **divergent**, otherwise.

... is a divergent rewrite sequence of length ω .

$$\blacksquare R = \{ f(x, x) \rightarrow f(a, b), a \rightarrow c(a), b \rightarrow c(b) \}$$

$$f(a, b) \rightarrow^\omega f(c^\omega, b) \rightarrow^\omega f(c^\omega, c^\omega) \rightarrow f(a, b)$$

... is a strongly convergent rewrite sequence of length $\omega \cdot 2 + 1$.

Definition

A reduction of length α is **strongly convergent** if for every limit ordinal $\lambda \leq \alpha$ the depth d_β tends to infinity as β approaches λ from below, and **divergent**, otherwise.

Example

$$1 \quad R = \{ a \rightarrow b, b \rightarrow a \}$$

$$a \rightarrow b \rightarrow a \rightarrow b \rightarrow \dots$$

Definition

A reduction of length α is **strongly convergent** if for every limit ordinal $\lambda \leq \alpha$ the depth d_β tends to infinity as β approaches λ from below, and **divergent**, otherwise.

Example

$$1 \quad R = \{ a \rightarrow b, b \rightarrow a \}$$

$$a \rightarrow b \rightarrow a \rightarrow b \rightarrow \dots$$

\dots is a divergent rewrite sequence of length ω .

Definition

A reduction of length α is **strongly convergent** if for every limit ordinal $\lambda \leq \alpha$ the depth d_β tends to infinity as β approaches λ from below, and **divergent**, otherwise.

Example

$$1 \quad R = \{ a \rightarrow b, b \rightarrow a \}$$

$$a \rightarrow b \rightarrow a \rightarrow b \rightarrow \dots$$

\dots is a divergent rewrite sequence of length ω .

$$2 \quad R = \{ f(x, x) \rightarrow f(a, b), a \rightarrow c(a), b \rightarrow c(b) \}$$

$$f(a, b) \rightarrow^\omega f(c^\omega, b) \rightarrow^\omega f(c^\omega, c^\omega) \rightarrow f(a, b)$$

Definition

A reduction of length α is **strongly convergent** if for every limit ordinal $\lambda \leq \alpha$ the depth d_β tends to infinity as β approaches λ from below, and **divergent**, otherwise.

Example

$$1 \quad R = \{ a \rightarrow b, b \rightarrow a \}$$

$$a \rightarrow b \rightarrow a \rightarrow b \rightarrow \dots$$

... is a divergent rewrite sequence of length ω .

$$2 \quad R = \{ f(x, x) \rightarrow f(a, b), a \rightarrow c(a), b \rightarrow c(b) \}$$

$$f(a, b) \rightarrow^\omega f(c^\omega, b) \rightarrow^\omega f(c^\omega, c^\omega) \rightarrow f(a, b)$$

... is a strongly convergent rewrite sequence of length $\omega \cdot 2 + 1$.

Definition

A reduction of length α is **strongly convergent** if for every limit ordinal $\lambda \leq \alpha$ the depth d_β tends to infinity as β approaches λ from below, and **divergent**, otherwise.

Example

$$\mathbf{1} \quad R = \{ a \rightarrow b, b \rightarrow a \}$$

$$a \rightarrow b \rightarrow a \rightarrow b \rightarrow \dots$$

... is a divergent rewrite sequence of length ω .

$$\mathbf{2} \quad R = \{ f(x, x) \rightarrow f(a, b), a \rightarrow c(a), b \rightarrow c(b) \}$$

$$f(a, b) \rightarrow^\omega f(c^\omega, b) \rightarrow^\omega f(c^\omega, c^\omega) \rightarrow f(a, b)$$

... is a strongly convergent rewrite sequence of length $\omega \cdot 2 + 1$.

Lemma

A reduction τ is strongly convergent

\iff for every $n \in \mathbb{N}$ there are only finitely many steps at depth n in τ .

Definition

We write $s \rightarrow\!\!\rightarrow t$ if the rewrite sequence is strongly convergent and with limit t .

But every prefix is convergent!

Definition

We write $s \twoheadrightarrow t$ if the rewrite sequence is strongly convergent and with limit t .

Example

$R = \{ a \rightarrow c(a) \}$. Then $a \twoheadrightarrow c^\omega$.

But every prefix is convergent!

Definition

We write $s \twoheadrightarrow t$ if the rewrite sequence is strongly convergent and with limit t .

Example

$R = \{ a \rightarrow c(a) \}$. Then $a \twoheadrightarrow c^\omega$.

Lemma

Every proper prefix of a (even divergent) rewrite sequence is strongly convergent.

But every prefix is convergent!

Definition

We write $s \twoheadrightarrow t$ if the rewrite sequence is strongly convergent and with limit t .

Example

$R = \{ a \rightarrow c(a) \}$. Then $a \twoheadrightarrow c^\omega$.

Lemma

Every proper prefix of a (even divergent) rewrite sequence is strongly convergent.

Example

$R = \{ f(x, x) \rightarrow f(a, b), a \rightarrow c(a), b \rightarrow c(b) \}$

$$f(a, b) \rightarrow^{\omega \cdot 2 + 1} f(a, b) \rightarrow^{\omega \cdot 2 + 1} f(a, b) \rightarrow^{\omega \cdot 2 + 1} \dots$$

... is a divergent rewrite sequence of length ω^2 .

Definition

We write $s \twoheadrightarrow t$ if the rewrite sequence is strongly convergent and with limit t .

Example

$R = \{ a \rightarrow c(a) \}$. Then $a \twoheadrightarrow c^\omega$.

Lemma

Every proper prefix of a (even divergent) rewrite sequence is strongly convergent.

Example

$R = \{ f(x, x) \rightarrow f(a, b), a \rightarrow c(a), b \rightarrow c(b) \}$

$$f(a, b) \rightarrow^{\omega \cdot 2 + 1} f(a, b) \rightarrow^{\omega \cdot 2 + 1} f(a, b) \rightarrow^{\omega \cdot 2 + 1} \dots$$

... is a divergent rewrite sequence of length ω^2 . But every prefix is convergent!

Comparison finitary vs. infinitary rewriting

finitary rewriting	infinitary rewriting
finite reduction	strongly convergent reduction
infinite reduction	divergent reduction

Then R is WN^∞ , SN^∞ and CR^∞ .

- Let $R = \{ a \rightarrow a, a \rightarrow c(a) \}$. Then R is WN^∞ and CR^∞ , but not SN^∞ .

Comparison finitary vs. infinitary rewriting

finitary rewriting	infinitary rewriting
finite reduction	strongly convergent reduction
infinite reduction	divergent reduction

Definition

Let \mathcal{R} be a TRS and s a term. Then the term s is

- **infinitary strongly normalizing** (SN^∞) if s admits no divergent reductions,
- **infinitary weakly normalizing** (WN^∞) if s admits a reduction to normal form,
- **infinitary confluent** (CR^∞) if $\forall t_1 \leftarrow\leftarrow s \rightarrow\rightarrow t_2. t_1 \rightarrow\rightarrow \cdot \leftarrow\leftarrow t_2$.

Likewise \mathcal{R} has the respective property if all terms from $\mathcal{T}^\infty(\Sigma, \mathcal{X})$ have.

Then R is WN^∞ , SN^∞ and CR^∞ .

- Let $R = \{ a \rightarrow a, a \rightarrow c(a) \}$. Then R is WN^∞ and CR^∞ , but not SN^∞ .

Comparison finitary vs. infinitary rewriting

finitary rewriting	infinitary rewriting
finite reduction	strongly convergent reduction
infinite reduction	divergent reduction

Definition

Let \mathcal{R} be a TRS and s a term. Then the term s is

- **infinitary strongly normalizing** (SN^∞) if s admits no divergent reductions,
- **infinitary weakly normalizing** (WN^∞) if s admits a reduction to normal form,
- **infinitary confluent** (CR^∞) if $\forall t_1 \leftarrow\leftarrow s \rightarrow\rightarrow t_2. t_1 \rightarrow\rightarrow \cdot \leftarrow\leftarrow t_2$.

Likewise \mathcal{R} has the respective property if all terms from $\mathcal{T}^\infty(\Sigma, \mathcal{X})$ have.

Example

- Let $R = \{ a \rightarrow c(a) \}$.

Comparison finitary vs. infinitary rewriting

finitary rewriting	infinitary rewriting
finite reduction	strongly convergent reduction
infinite reduction	divergent reduction

Definition

Let \mathcal{R} be a TRS and s a term. Then the term s is

- **infinitary strongly normalizing** (SN^∞) if s admits no divergent reductions,
- **infinitary weakly normalizing** (WN^∞) if s admits a reduction to normal form,
- **infinitary confluent** (CR^∞) if $\forall t_1 \leftarrow\!\!\leftarrow s \twoheadrightarrow\!\!\rightarrow t_2. t_1 \twoheadrightarrow\!\!\rightarrow \cdot \leftarrow\!\!\leftarrow t_2$.

Likewise \mathcal{R} has the respective property if all terms from $\mathcal{T}^\infty(\Sigma, \mathcal{X})$ have.

Example

- Let $R = \{ a \rightarrow c(a) \}$. Then R is WN^∞ , SN^∞ and CR^∞ .

Comparison finitary vs. infinitary rewriting

finitary rewriting	infinitary rewriting
finite reduction	strongly convergent reduction
infinite reduction	divergent reduction

Definition

Let \mathcal{R} be a TRS and s a term. Then the term s is

- **infinitary strongly normalizing** (SN^∞) if s admits no divergent reductions,
- **infinitary weakly normalizing** (WN^∞) if s admits a reduction to normal form,
- **infinitary confluent** (CR^∞) if $\forall t_1 \leftarrow\leftarrow s \rightarrow\rightarrow t_2. t_1 \rightarrow\rightarrow \cdot \leftarrow\leftarrow t_2$.

Likewise \mathcal{R} has the respective property if all terms from $\mathcal{T}^\infty(\Sigma, \mathcal{X})$ have.

Example

- Let $R = \{ a \rightarrow c(a) \}$. Then R is WN^∞ , SN^∞ and CR^∞ .
- Let $R = \{ a \rightarrow a, a \rightarrow c(a) \}$.

Comparison finitary vs. infinitary rewriting

finitary rewriting	infinitary rewriting
finite reduction	strongly convergent reduction
infinite reduction	divergent reduction

Definition

Let \mathcal{R} be a TRS and s a term. Then the term s is

- **infinitary strongly normalizing** (SN^∞) if s admits no divergent reductions,
- **infinitary weakly normalizing** (WN^∞) if s admits a reduction to normal form,
- **infinitary confluent** (CR^∞) if $\forall t_1 \leftarrow\!\!\leftarrow s \twoheadrightarrow\!\!\twoheadrightarrow t_2. t_1 \twoheadrightarrow\!\!\twoheadrightarrow \cdot \leftarrow\!\!\leftarrow t_2$.

Likewise \mathcal{R} has the respective property if all terms from $\mathcal{T}^\infty(\Sigma, \mathcal{X})$ have.

Example

- Let $R = \{ a \rightarrow c(a) \}$. Then R is WN^∞ , SN^∞ and CR^∞ .
- Let $R = \{ a \rightarrow a, a \rightarrow c(a) \}$. Then R is WN^∞ and CR^∞ , but not SN^∞ .

Remark

- $SN^\infty \not\Rightarrow SN$

$$a \rightarrow c(a)$$

Here, $a \rightarrow^* c^\omega$ which is a normal form.

Even one collapsing rule is sufficient to violate CR^∞ .

Take $R = \{ f(x, y) \rightarrow y \}$. Then

$$f(x, f(x, f(x, \dots))) \leftarrow f(x, f(y, f(x, f(y, \dots)))) \rightarrow f(y, f(y, f(y, \dots)))$$

Remark

- $SN^\infty \not\Rightarrow SN \vee WN$

$$a \rightarrow c(a)$$

Here, $a \rightarrow^* c^\omega$ which is a normal form.

Even one collapsing rule is sufficient to violate CR^∞ .

Take $R = \{ f(x, y) \rightarrow y \}$. Then

$$f(x, f(x, f(x, \dots))) \leftarrow f(x, f(y, f(x, f(y, \dots)))) \rightarrow f(y, f(y, f(y, \dots)))$$

Remark

- $SN^\infty \not\Rightarrow SN \vee WN$

$$a \rightarrow c(a)$$

Here, $a \rightarrow^* c^\omega$ which is a normal form.

- $SN \not\Rightarrow SN^\infty$

Even one collapsing rule is sufficient to violate CR^∞ .

Take $R = \{ f(x, y) \rightarrow y \}$. Then

$$f(x, f(x, f(x, \dots))) \leftarrow f(x, f(y, f(x, f(y, \dots)))) \rightarrow f(y, f(y, f(y, \dots)))$$

Remark

- $SN^\infty \not\Rightarrow SN \vee WN$

$$a \rightarrow c(a)$$

Here, $a \rightarrow^* c^\omega$ which is a normal form.

- $SN \not\Rightarrow SN^\infty$

$$I(x) \rightarrow x$$

Here, $I(I(I(\dots)))$ rewrites only to itself.

Even one collapsing rule is sufficient to violate CR^∞ .

Take $R = \{ f(x, y) \rightarrow y \}$. Then

$$f(x, f(x, f(x, \dots))) \leftarrow f(x, f(y, f(x, f(y, \dots)))) \rightarrow f(y, f(y, f(y, \dots)))$$

Remark

- $SN^\infty \not\Rightarrow SN \vee WN$

$$a \rightarrow c(a)$$

Here, $a \rightarrow^* c^\omega$ which is a normal form.

- $SN \not\Rightarrow SN^\infty \vee WN^\infty$

$$I(x) \rightarrow x$$

Here, $I(I(I(\dots)))$ rewrites only to itself.

Even one collapsing rule is sufficient to violate CR^∞ .

Take $R = \{ f(x, y) \rightarrow y \}$. Then

$$f(x, f(x, f(x, \dots))) \leftarrow f(x, f(y, f(x, f(y, \dots)))) \rightarrow f(y, f(y, f(y, \dots)))$$

Remark

- $SN^\infty \not\Rightarrow SN \vee WN$

$$a \rightarrow c(a)$$

Here, $a \rightarrow^* c^\omega$ which is a normal form.

- $SN \not\Rightarrow SN^\infty \vee WN^\infty$

$$I(x) \rightarrow x$$

Here, $I(I(I(\dots)))$ rewrites only to itself.

- $CR^\infty \quad CR$

Even one collapsing rule is sufficient to violate CR^∞ .

Take $R = \{ f(x, y) \rightarrow y \}$. Then

$$f(x, f(x, f(x, \dots))) \leftarrow f(x, f(y, f(x, f(y, \dots)))) \rightarrow^* f(y, f(y, f(y, \dots)))$$

Remark

- $SN^\infty \not\Rightarrow SN \vee WN$

$$a \rightarrow c(a)$$

Here, $a \rightarrow^* c^\omega$ which is a normal form.

- $SN \not\Rightarrow SN^\infty \vee WN^\infty$

$$I(x) \rightarrow x$$

Here, $I(I(I(\dots)))$ rewrites only to itself.

- $CR^\infty \not\Rightarrow CR$

$$a \rightarrow b, a \rightarrow c, b \rightarrow d(b), c \rightarrow d(c)$$

Here, $\neg(b \downarrow c)$, but $b \rightarrow^* d^\omega \leftarrow c$.

Even one collapsing rule is sufficient to violate CR^∞ .

Take $R = \{ f(x, y) \rightarrow y \}$. Then

$$f(x, f(x, f(x, \dots))) \leftarrow f(x, f(y, f(x, f(y, \dots)))) \rightarrow^* f(y, f(y, f(y, \dots)))$$

Remark

- $SN^\infty \not\Rightarrow SN \vee WN$

$$a \rightarrow c(a)$$

Here, $a \rightarrow^* c^\omega$ which is a normal form.

- $SN \not\Rightarrow SN^\infty \vee WN^\infty$

$$I(x) \rightarrow x$$

Here, $I(I(I(\dots)))$ rewrites only to itself.

- $CR^\infty \not\Rightarrow CR$

$$a \rightarrow b, a \rightarrow c, b \rightarrow d(b), c \rightarrow d(c)$$

Here, $\neg(b \downarrow c)$, but $b \rightarrow^* d^\omega \leftarrow c$.

- $CR \not\Rightarrow CR^\infty$

Even one collapsing rule is sufficient to violate CR^∞ .

Take $R = \{ f(x, y) \rightarrow y \}$. Then

$$f(x, f(x, f(x, \dots))) \leftarrow f(x, f(y, f(x, f(y, \dots)))) \rightarrow^* f(y, f(y, f(y, \dots)))$$

Remark

- $SN^\infty \not\Rightarrow SN \vee WN$

$$a \rightarrow c(a)$$

Here, $a \rightarrow^* c^\omega$ which is a normal form.

- $SN \not\Rightarrow SN^\infty \vee WN^\infty$

$$I(x) \rightarrow x$$

Here, $I(I(I(\dots)))$ rewrites only to itself.

- $CR^\infty \not\Rightarrow CR$

$$a \rightarrow b, a \rightarrow c, b \rightarrow d(b), c \rightarrow d(c)$$

Here, $\neg(b \downarrow c)$, but $b \rightarrow^* d^\omega \leftarrow c$.

- $CR \not\Rightarrow CR^\infty$

$$A(x) \rightarrow x, B(x) \rightarrow x$$

Here, $A^\omega \leftarrow (AB)^\omega \rightarrow^* B^\omega$.

Even one collapsing rule is sufficient to violate CR^∞ .

Take $R = \{ f(x, y) \rightarrow y \}$. Then

$$f(x, f(x, f(x, \dots))) \leftarrow f(x, f(y, f(x, f(y, \dots)))) \rightarrow^* f(y, f(y, f(y, \dots)))$$

Remark

- $SN^\infty \not\Rightarrow SN \vee WN$

$$a \rightarrow c(a)$$

Here, $a \rightarrow^{\omega} c^{\omega}$ which is a normal form.

- $SN \not\Rightarrow SN^\infty \vee WN^\infty$

$$I(x) \rightarrow x$$

Here, $I(I(I(\dots)))$ rewrites only to itself.

- $CR^\infty \not\Rightarrow CR$

$$a \rightarrow b, a \rightarrow c, b \rightarrow d(b), c \rightarrow d(c)$$

Here, $\neg(b \downarrow c)$, but $b \rightarrow^{\omega} d^{\omega} \leftarrow c$.

- $CR \not\Rightarrow CR^\infty$

$$A(x) \rightarrow x, B(x) \rightarrow x$$

Here, $A^{\omega} \leftarrow (AB)^{\omega} \rightarrow B^{\omega}$.

Remark

The example $A(x) \rightarrow x, B(x) \rightarrow x$ shows: not every orthogonal TRSs is CR^∞ .

Remark

- $SN^\infty \not\Rightarrow SN \vee WN$
 $a \rightarrow c(a)$
Here, $a \rightarrow^{\omega} c^{\omega}$ which is a normal form.
- $SN \not\Rightarrow SN^\infty \vee WN^\infty$
 $I(x) \rightarrow x$
Here, $I(I(I(\dots)))$ rewrites only to itself.
- $CR^\infty \not\Rightarrow CR$
 $a \rightarrow b, a \rightarrow c, b \rightarrow d(b), c \rightarrow d(c)$
Here, $\neg(b \downarrow c)$, but $b \rightarrow^{\omega} d^{\omega} \leftarrow c$.
- $CR \not\Rightarrow CR^\infty$
 $A(x) \rightarrow x, B(x) \rightarrow x$
Here, $A^{\omega} \leftarrow (AB)^{\omega} \rightarrow B^{\omega}$.

Remark

The example $A(x) \rightarrow x, B(x) \rightarrow x$ shows: not every orthogonal TRSs is CR^∞ .
 Even one collapsing rule is sufficient to violate CR^∞ .

Remark

- $SN^\infty \not\Rightarrow SN \vee WN$
 $a \rightarrow c(a)$
Here, $a \rightarrow^ c^\omega$ which is a normal form.*
- $SN \not\Rightarrow SN^\infty \vee WN^\infty$
 $I(x) \rightarrow x$
Here, $I(I(I(\dots)))$ rewrites only to itself.
- $CR^\infty \not\Rightarrow CR$
 $a \rightarrow b, a \rightarrow c, b \rightarrow d(b), c \rightarrow d(c)$
Here, $\neg(b \downarrow c)$, but $b \rightarrow^ d^\omega \leftarrow c$.*
- $CR \not\Rightarrow CR^\infty$
 $A(x) \rightarrow x, B(x) \rightarrow x$
Here, $A^\omega \leftarrow (AB)^\omega \rightarrow^ B^\omega$.*

Remark

The example $A(x) \rightarrow x, B(x) \rightarrow x$ shows: not every orthogonal TRSs is CR^∞ .

Even one collapsing rule is sufficient to violate CR^∞ .

Take $R = \{ f(x, y) \rightarrow y \}$. Then

$$f(x, f(x, f(x, \dots))) \leftarrow f(x, f(y, f(x, f(y, \dots)))) \rightarrow^* f(y, f(y, f(y, \dots)))$$

Remark (The failure of Newmann's Lemma for infinitary rewriting)

$$WCR \wedge SN^\infty \not\Rightarrow CR^\infty$$

Remark (The failure of Newmann's Lemma for infinitary rewriting)

$$WCR \wedge SN^\infty \not\Rightarrow CR^\infty$$

For example:

$$R = \{ a \rightarrow b(a), \\ a \rightarrow c(a), \\ c(b(x)) \rightarrow b(b(x)) \}$$

is WCR and SN^∞ , but not CR^∞ .

Results for (Weakly) Orthogonal TRSs

Theorem

Every weakly orthogonal TRS *without collapsing rules* is CR^∞ .

Results for (Weakly) Orthogonal TRSs

Theorem

Every weakly orthogonal TRS *without collapsing rules* is CR^∞ .

Definition

A TRS \mathcal{R} is UN^∞ if $s \llcorner \cdot \rhd t \Rightarrow s = t$ for all normal forms $s, t \in \mathcal{T}^\infty(\Sigma, \mathcal{X})$.

Results for (Weakly) Orthogonal TRSs

Theorem

Every weakly orthogonal TRS *without collapsing rules* is CR^∞ .

Definition

A TRS \mathcal{R} is UN^∞ if $s \leftarrow\!\!\leftarrow \cdot \rightarrow\!\!\rightarrow t \Rightarrow s = t$ for all normal forms $s, t \in \mathcal{T}^\infty(\Sigma, \mathcal{X})$.

Theorem

Every orthogonal TRS is UN^∞ .

Results for (Weakly) Orthogonal TRSs

Theorem

Every weakly orthogonal TRS *without collapsing rules* is CR^∞ .

Definition

A TRS \mathcal{R} is UN^∞ if $s \leftarrow \cdot \rightarrow t \Rightarrow s = t$ for all normal forms $s, t \in \mathcal{T}^\infty(\Sigma, \mathcal{X})$.

Theorem

Every orthogonal TRS is UN^∞ .

Example

Weakly orthogonal TRSs are not necessarily UN^∞ :

$$S(P(x)) \rightarrow x$$

$$P(S(x)) \rightarrow x$$

Then

$$S^\omega \leftarrow S^1(P^2(S^3(P^4(\dots)))) \rightarrow P^\omega$$

Compression and Parallel Moves

Theorem (Compression)

Let \mathcal{R} be an left-linear TRS. Then $s \rightarrow^ t$ implies $s \rightarrow^{\leq \omega} t$.*

That is, every strongly convergent reduction can be compressed to length $\leq \omega$.

Compression and Parallel Moves

Theorem (Compression)

Let \mathcal{R} be an left-linear TRS. Then $s \twoheadrightarrow t$ implies $s \rightarrow^{\leq \omega} t$.

That is, every strongly convergent reduction can be compressed to length $\leq \omega$.

Theorem (Parallel Moves)

Let \mathcal{R} be an orthogonal TRS. Then $t_1 \ll s \twoheadrightarrow t_2 \Rightarrow t_1 \twoheadrightarrow \cdot \ll t_2$.

