- Lecture 1: Introduction, Abstract Rewriting
- Lecture 2: Term Rewriting
- Lecture 3: Combinatory Logic
- Lecture 4: Termination
- Lecture 5: Matching, Unification
- Lecture 6: Equational Reasoning, Completion
- Lecture 7: Confluence
- Lecture 8: Modularity
- Lecture 9: Strategies
- Lecture 10: Decidability
- Lecture 11: Infinitary Rewriting

Outline

- Overview
- Strategies

Strategies

Definition

A rewrite strategy $\mathcal{S}$ is mapping that assigns to every reducible term t a nonempty set of finite nonempty rewrite sequences starting from t.

Definition

A rewrite strategy $\mathcal{S}$ is mapping that assigns to every reducible term t a nonempty set of finite nonempty rewrite sequences starting from t.

- $\mathcal{S}$ is deterministic if $|\mathcal{S}(t)| = 1$ for every reducible term t

Definition

A rewrite strategy $\mathcal{S}$ is mapping that assigns to every reducible term t a nonempty set of finite nonempty rewrite sequences starting from t.

- $\mathcal{S}$ is deterministic if $|\mathcal{S}(t)| = 1$ for every reducible term t
- $\mathcal{S}$ normalizes term t if there are no infinite $\mathcal{S}$ rewrite sequences starting from t

Definition

A rewrite strategy $\mathcal{S}$ is mapping that assigns to every reducible term t a nonempty set of finite nonempty rewrite sequences starting from t.

- $\mathcal{S}$ is deterministic if $|\mathcal{S}(t)| = 1$ for every reducible term t
- $\mathcal{S}$ normalizes term t if there are no infinite $\mathcal{S}$ rewrite sequences starting from t
- $\mathcal{S}$ is normalizing if it normalizes every term that has a normal form

Definition

A rewrite strategy $\mathcal{S}$ is mapping that assigns to every reducible term t a nonempty set of finite nonempty rewrite sequences starting from t.

- $\mathcal{S}$ is deterministic if $|\mathcal{S}(t)| = 1$ for every reducible term t
- $\mathcal{S}$ normalizes term t if there are no infinite $\mathcal{S}$ rewrite sequences starting from t
- $\mathcal{S}$ is normalizing if it normalizes every term that has a normal form
- $\mathcal{S}$ is perpetual if every maximal $\mathcal{S}$ rewrite sequence starting from any non-terminating term is infinite

Definition

A rewrite sequence is maximal if it is infinite, or it ends in a normal form.

Definition

A rewrite strategy $\mathcal{S}$ is mapping that assigns to every reducible term t a nonempty set of finite nonempty rewrite sequences starting from t.

- $\mathcal{S}$ is deterministic if $|\mathcal{S}(t)| = 1$ for every reducible term t
- $\mathcal{S}$ normalizes term t if there are no infinite $\mathcal{S}$ rewrite sequences starting from t
- $\mathcal{S}$ is normalizing if it normalizes every term that has a normal form
- $\mathcal{S}$ is perpetual if every maximal $\mathcal{S}$ rewrite sequence starting from any non-terminating term is infinite

Definition

A rewrite sequence is maximal if it is infinite, or it ends in a normal form.

Lemma

For terminating TRSs every strategy is normalizing and perpetual.

Definition

Let $\rho = t_0 \to t_1 \to \ldots$ be a finite or infinite rewrite sequence.

Definition

Let $\rho = t_0 \to t_1 \to \ldots$ be a finite or infinite rewrite sequence.

- Consider a redex occurrence s is some term t_n of ρ.

Definition

Let $\rho = t_0 \to t_1 \to \ldots$ be a finite or infinite rewrite sequence.

- Consider a redex occurrence s is some term t_n of ρ.

  Then s is secured if eventually there are no residuals of s left.

Definition

Let $\rho = t_0 \to t_1 \to \ldots$ be a finite or infinite rewrite sequence.

- Consider a redex occurrence s is some term t_n of ρ.

  Then s is secured if eventually there are no residuals of s left.

  That is, there exits $m > n$ such that t_m contains no residuals of s.

Definition

Let $\rho = t_0 \to t_1 \to \ldots$ be a finite or infinite rewrite sequence.

- Consider a redex occurrence s is some term t_n of ρ.

  Then s is secured if eventually there are no residuals of s left.

  That is, there exits $m > n$ such that t_m contains no residuals of s.

- The reduction ρ is fair if every redex occurring in ρ is secured.

Definition

Let $\rho = t_0 \to t_1 \to \ldots$ be a finite or infinite rewrite sequence.

- Consider a redex occurrence s is some term t_n of ρ.

  Then s is secured if eventually there are no residuals of s left.

  That is, there exits $m > n$ such that t_m contains no residuals of s.

- The reduction ρ is fair if every redex occurring in ρ is secured.

Definition

A strategy $\mathcal{S}$ is fair if every every maximal $\mathcal{S}$ rewrite sequence is fair.

Definition

A one-step strategy maps every reducible term to a set of one-step reductions.

Example

There exists no fair one-step strategy for $\mathcal{R} = \{I(x) \to I(x)\}$.

Definition

A one-step strategy maps every reducible term to a set of one-step reductions.

Example

There exists no fair one-step strategy for $\mathcal{R} = \{I(x) \to I(x)\}$.

For the term:

$$t = I(I(x))$$

there are only 3 possible mappings:

Definition

A one-step strategy maps every reducible term to a set of one-step reductions.

Example

There exists no fair one-step strategy for $\mathcal{R} = \{I(x) \to I(x)\}$.

For the term:

$$t = I(I(x))$$

there are only 3 possible mappings:

- $\mathcal{S}(t) = \{I(I(x)) \to_\varepsilon I(I(x))\}$,

Definition

A one-step strategy maps every reducible term to a set of one-step reductions.

Example

There exists no fair one-step strategy for $\mathcal{R} = \{I(x) \to I(x)\}$.

For the term:

$$t = I(I(x))$$

there are only 3 possible mappings:

- $\mathcal{S}(t) = \{I(I(x)) \to_\varepsilon I(I(x))\}$,
- $\mathcal{S}(t) = \{I(I(x)) \to_1 I(I(x))\}$, or

Definition

A one-step strategy maps every reducible term to a set of one-step reductions.

Example

There exists no fair one-step strategy for $\mathcal{R} = \{I(x) \to I(x)\}$.

For the term:

$$t = I(I(x))$$

there are only 3 possible mappings:

- $\mathcal{S}(t) = \{I(I(x)) \to_\varepsilon I(I(x))\}$,
- $\mathcal{S}(t) = \{I(I(x)) \to_1 I(I(x))\}$, or
- $\mathcal{S}(t) = \{I(I(x)) \to_\varepsilon I(I(x)),\ I(I(x)) \to_1 I(I(x))\}$.

Definition

A one-step strategy maps every reducible term to a set of one-step reductions.

Example

There exists no fair one-step strategy for $\mathcal{R} = \{I(x) \to I(x)\}$.

For the term:

$$t = I(I(x))$$

there are only 3 possible mappings:

- $\mathcal{S}(t) = \{I(I(x)) \to_\varepsilon I(I(x))\}$,
- $\mathcal{S}(t) = \{I(I(x)) \to_1 I(I(x))\}$, or
- $\mathcal{S}(t) = \{I(I(x)) \to_\varepsilon I(I(x)),\ I(I(x)) \to_1 I(I(x))\}$.

None of these is fair as we can always continue to reduce the same occurrence of I.

Example

- rewrite rules

$$
\begin{array}{llll}
0+0 \to 0 & 1+0 \to 1 & \cdots & 9+0 \to 9 \\
0+1 \to 1 & 1+1 \to 2 & \cdots & 9+1 \to 1:0 \\
0+2 \to 2 & 1+2 \to 3 & \cdots & 9+2 \to 1:1 \\
0+3 \to 3 & 1+3 \to 4 & \cdots & 9+3 \to 1:2 \\
0+4 \to 4 & 1+4 \to 5 & \cdots & 9+4 \to 1:3 \\
0+5 \to 5 & 1+5 \to 6 & \cdots & 9+5 \to 1:4 \\
0+6 \to 6 & 1+6 \to 7 & \cdots & 9+6 \to 1:5 \\
0+7 \to 7 & 1+7 \to 8 & \cdots & 9+7 \to 1:6 \\
0+8 \to 8 & 1+8 \to 9 & \cdots & 9+8 \to 1:7 \\
0+9 \to 9 & 1+9 \to 1:0 & \cdots & 9+9 \to 1:8
\end{array}
$$

$$x + (y : z) \to y : (x + z) \qquad 0 : x \to x$$

$$(x : y) + z \to x : (y + z) \qquad x : (y : z) \to (x + y) : z$$

- term

$$((0 : (1 + 2)) + (3 + 4)) + (5 + 6)$$

Example (cont'd)

term

$$\boxed{\boxed{0 : \boxed{1 + 2}} + \boxed{3 + 4}} + \boxed{5 + 6}$$

tree representation



Example (cont'd)

term

$$\boxed{\boxed{0 : \boxed{1 + 2}} + \boxed{3 + 4}} + \boxed{5 + 6}$$

tree representation



outermost redexes

Example (cont'd)

term

$$\boxed{\boxed{0 : \boxed{1 + 2}} + \boxed{3 + 4}} + \boxed{5 + 6}$$

tree representation



innermost redexes

Example (cont'd)

term

$$\boxed{\boxed{0 : \boxed{1 + 2}} + \boxed{3 + 4}} + \boxed{5 + 6}$$

tree representation



leftmost outermost strategy

Example (cont'd)

term

$$\boxed{\boxed{0 : \boxed{1 + 2}} + \boxed{3 + 4}} + \boxed{5 + 6}$$

tree representation



leftmost innermost strategy

Example (cont'd)

term

$$\boxed{\boxed{0 : \boxed{1 + 2}} + \boxed{3 + 4}} + \boxed{5 + 6}$$

tree representation



parallel outermost strategy

Example (cont'd)

term

$$\boxed{\boxed{0 : \boxed{1 + 2}} + \boxed{3 + 4}} + \boxed{5 + 6}$$

tree representation



parallel innermost strategy

Definition

Leftmost outermost strategy always reduces the leftmost of the outermost redexes.

Definition

Leftmost outermost strategy always reduces the leftmost of the outermost redexes.

Example (cont'd)

$$((0 : (1 + 2)) + (3 + 4)) + (5 + 6)$$

Definition

Leftmost outermost strategy always reduces the leftmost of the outermost redexes.

Example (cont'd)

$$((0 : (1 + 2)) + (3 + 4)) + (5 + 6) \rightarrow (0 : ((1 + 2) + (3 + 4))) + (5 + 6)$$

Definition

Leftmost outermost strategy always reduces the leftmost of the outermost redexes.

Example (cont'd)

$$((0 : (1 + 2)) + (3 + 4)) + (5 + 6) \rightarrow (0 : ((1 + 2) + (3 + 4))) + (5 + 6)$$
$$\rightarrow 0 : (((1 + 2) + (3 + 4)) + (5 + 6))$$

Definition

Leftmost outermost strategy always reduces the leftmost of the outermost redexes.

Example (cont'd)

$$((0 : (1 + 2)) + (3 + 4)) + (5 + 6) \rightarrow (0 : ((1 + 2) + (3 + 4))) + (5 + 6)$$
$$\rightarrow 0 : (((1 + 2) + (3 + 4)) + (5 + 6))$$
$$\rightarrow ((1 + 2) + (3 + 4)) + (5 + 6)$$

Definition

Leftmost outermost strategy always reduces the leftmost of the outermost redexes.

Example (cont'd)

$$\begin{aligned}((0 : (1 + 2)) + (3 + 4)) + (5 + 6) &\to (0 : ((1 + 2) + (3 + 4))) + (5 + 6) \\ &\to 0 : (((1 + 2) + (3 + 4)) + (5 + 6)) \\ &\to ((1 + 2) + (3 + 4)) + (5 + 6) \\ &\to (3 + (3 + 4)) + (5 + 6)\end{aligned}$$

Definition

Leftmost outermost strategy always reduces the leftmost of the outermost redexes.

Example (cont'd)

$$\begin{aligned}((0 : (1 + 2)) + (3 + 4)) + (5 + 6) &\to (0 : ((1 + 2) + (3 + 4))) + (5 + 6) \\ &\to 0 : (((1 + 2) + (3 + 4)) + (5 + 6)) \\ &\to ((1 + 2) + (3 + 4)) + (5 + 6) \\ &\to (3 + (3 + 4)) + (5 + 6) \\ &\to (3 + 7) + (5 + 6)\end{aligned}$$

Definition

Leftmost outermost strategy always reduces the leftmost of the outermost redexes.

Example (cont'd)

$$\begin{aligned}((0 : (1 + 2)) + (3 + 4)) + (5 + 6) &\to (0 : ((1 + 2) + (3 + 4))) + (5 + 6) \\ &\to 0 : (((1 + 2) + (3 + 4)) + (5 + 6)) \\ &\to ((1 + 2) + (3 + 4)) + (5 + 6) \\ &\to (3 + (3 + 4)) + (5 + 6) \\ &\to (3 + 7) + (5 + 6) \\ &\to (1 : 0) + (5 + 6)\end{aligned}$$

Definition

Leftmost outermost strategy always reduces the leftmost of the outermost redexes.

Example (cont'd)

$$\begin{aligned}((0 : (1 + 2)) + (3 + 4)) + (5 + 6) &\to (0 : ((1 + 2) + (3 + 4))) + (5 + 6)\\ &\to 0 : (((1 + 2) + (3 + 4)) + (5 + 6))\\ &\to ((1 + 2) + (3 + 4)) + (5 + 6)\\ &\to (3 + (3 + 4)) + (5 + 6)\\ &\to (3 + 7) + (5 + 6)\\ &\to (1 : 0) + (5 + 6)\\ &\to 1 : (0 + (5 + 6))\end{aligned}$$

Definition

Leftmost outermost strategy always reduces the leftmost of the outermost redexes.

Example (cont'd)

$$
\begin{aligned}
((0 : (1 + 2)) + (3 + 4)) + (5 + 6) &\to (0 : ((1 + 2) + (3 + 4))) + (5 + 6) \\
&\to 0 : (((1 + 2) + (3 + 4)) + (5 + 6)) \\
&\to ((1 + 2) + (3 + 4)) + (5 + 6) \\
&\to (3 + (3 + 4)) + (5 + 6) \\
&\to (3 + 7) + (5 + 6) \\
&\to (1 : 0) + (5 + 6) \\
&\to 1 : (0 + (5 + 6)) \\
&\to 1 : (0 + (1 : 1))
\end{aligned}
$$

Definition

Leftmost outermost strategy always reduces the leftmost of the outermost redexes.

Example (cont'd)

$$\begin{aligned}((0 : (1 + 2)) + (3 + 4)) + (5 + 6) &\to (0 : ((1 + 2) + (3 + 4))) + (5 + 6)\\ &\to 0 : (((1 + 2) + (3 + 4)) + (5 + 6))\\ &\to ((1 + 2) + (3 + 4)) + (5 + 6)\\ &\to (3 + (3 + 4)) + (5 + 6)\\ &\to (3 + 7) + (5 + 6)\\ &\to (1 : 0) + (5 + 6)\\ &\to 1 : (0 + (5 + 6))\\ &\to 1 : (0 + (1 : 1))\\ &\to 1 : (1 : (0 + 1))\end{aligned}$$

Definition

Leftmost outermost strategy always reduces the leftmost of the outermost redexes.

Example (cont'd)

$$
\begin{aligned}
((0 : (1 + 2)) + (3 + 4)) + (5 + 6) &\to (0 : ((1 + 2) + (3 + 4))) + (5 + 6) \\
&\to 0 : (((1 + 2) + (3 + 4)) + (5 + 6)) \\
&\to ((1 + 2) + (3 + 4)) + (5 + 6) \\
&\to (3 + (3 + 4)) + (5 + 6) \\
&\to (3 + 7) + (5 + 6) \\
&\to (1 : 0) + (5 + 6) \\
&\to 1 : (0 + (5 + 6)) \\
&\to 1 : (0 + (1 : 1)) \\
&\to 1 : (1 : (0 + 1)) \\
&\to (1 + 1) : (0 + 1)
\end{aligned}
$$

Definition

Leftmost outermost strategy always reduces the leftmost of the outermost redexes.

Example (cont'd)

$$
\begin{aligned}
((0 : (1 + 2)) + (3 + 4)) + (5 + 6) &\to (0 : ((1 + 2) + (3 + 4))) + (5 + 6) \\
&\to 0 : (((1 + 2) + (3 + 4)) + (5 + 6)) \\
&\to ((1 + 2) + (3 + 4)) + (5 + 6) \\
&\to (3 + (3 + 4)) + (5 + 6) \\
&\to (3 + 7) + (5 + 6) \\
&\to (1 : 0) + (5 + 6) \\
&\to 1 : (0 + (5 + 6)) \\
&\to 1 : (0 + (1 : 1)) \\
&\to 1 : (1 : (0 + 1)) \\
&\to (1 + 1) : (0 + 1) \\
&\to 2 : (0 + 1)
\end{aligned}
$$

Definition

Leftmost outermost strategy always reduces the leftmost of the outermost redexes.

Example (cont'd)

$$
\begin{aligned}
((0 : (1 + 2)) + (3 + 4)) + (5 + 6) &\to (0 : ((1 + 2) + (3 + 4))) + (5 + 6) \\
&\to 0 : (((1 + 2) + (3 + 4)) + (5 + 6)) \\
&\to ((1 + 2) + (3 + 4)) + (5 + 6) \\
&\to (3 + (3 + 4)) + (5 + 6) \\
&\to (3 + 7) + (5 + 6) \\
&\to (1 : 0) + (5 + 6) \\
&\to 1 : (0 + (5 + 6)) \\
&\to 1 : (0 + (1 : 1)) \\
&\to 1 : (1 : (0 + 1)) \\
&\to (1 + 1) : (0 + 1) \\
&\to 2 : (0 + 1) \\
&\to 2 : 1
\end{aligned}
$$

Definition

Leftmost innermost strategy always reduces the leftmost of the innermost redexes.

Definition

Leftmost innermost strategy always reduces the leftmost of the innermost redexes.

Example (cont'd)

$$(0 : (1 + 2)) + (3 + 4) + (5 + 6)$$

Definition

Leftmost innermost strategy always reduces the leftmost of the innermost redexes.

Example (cont'd)

$$(0 : (1 + 2)) + (3 + 4) + (5 + 6) \rightarrow ((0 : 3) + (3 + 4)) + (5 + 6)$$

Definition

Leftmost innermost strategy always reduces the leftmost of the innermost redexes.

Example (cont'd)

$$(0 : (1 + 2)) + (3 + 4) + (5 + 6) \to ((0 : 3) + (3 + 4)) + (5 + 6)$$
$$\to (3 + (3 + 4)) + (5 + 6)$$

Definition

Leftmost innermost strategy always reduces the leftmost of the innermost redexes.

Example (cont'd)

$$\begin{aligned}(0 : (1 + 2)) + (3 + 4) + (5 + 6) &\to ((0 : 3) + (3 + 4)) + (5 + 6)\\ &\to (3 + (3 + 4)) + (5 + 6)\\ &\to (3 + 7) + (5 + 6)\end{aligned}$$

Definition

Leftmost innermost strategy always reduces the leftmost of the innermost redexes.

Example (cont'd)

$$
\begin{aligned}
(0 : (1 + 2)) + (3 + 4) + (5 + 6) &\to ((0 : 3) + (3 + 4)) + (5 + 6) \\
&\to (3 + (3 + 4)) + (5 + 6) \\
&\to (3 + 7) + (5 + 6) \\
&\to (1 : 0) + (5 + 6)
\end{aligned}
$$

Definition

Leftmost innermost strategy always reduces the leftmost of the innermost redexes.

Example (cont'd)

$$
\begin{aligned}
(0 : (1 + 2)) + (3 + 4) + (5 + 6) &\to ((0 : 3) + (3 + 4)) + (5 + 6) \\
&\to (3 + (3 + 4)) + (5 + 6) \\
&\to (3 + 7) + (5 + 6) \\
&\to (1 : 0) + (5 + 6) \\
&\to (1 : 0) + (1 : 1)
\end{aligned}
$$

Definition

Leftmost innermost strategy always reduces the leftmost of the innermost redexes.

Example (cont'd)

$$
\begin{aligned}
(0 : (1 + 2)) + (3 + 4) + (5 + 6) &\to ((0 : 3) + (3 + 4)) + (5 + 6) \\
&\to (3 + (3 + 4)) + (5 + 6) \\
&\to (3 + 7) + (5 + 6) \\
&\to (1 : 0) + (5 + 6) \\
&\to (1 : 0) + (1 : 1) \\
&\to 1 : (0 + (1 : 1))
\end{aligned}
$$

Definition

Leftmost innermost strategy always reduces the leftmost of the innermost redexes.

Example (cont'd)

$$
\begin{aligned}
(0 : (1 + 2)) + (3 + 4) + (5 + 6) &\to ((0 : 3) + (3 + 4)) + (5 + 6) \\
&\to (3 + (3 + 4)) + (5 + 6) \\
&\to (3 + 7) + (5 + 6) \\
&\to (1 : 0) + (5 + 6) \\
&\to (1 : 0) + (1 : 1) \\
&\to 1 : (0 + (1 : 1)) \\
&\to 1 : (1 : (0 + 1))
\end{aligned}
$$

Definition

Leftmost innermost strategy always reduces the leftmost of the innermost redexes.

Example (cont'd)

$$
\begin{aligned}
(0 : (1 + 2)) + (3 + 4) + (5 + 6) &\to ((0 : 3) + (3 + 4)) + (5 + 6) \\
&\to (3 + (3 + 4)) + (5 + 6) \\
&\to (3 + 7) + (5 + 6) \\
&\to (1 : 0) + (5 + 6) \\
&\to (1 : 0) + (1 : 1) \\
&\to 1 : (0 + (1 : 1)) \\
&\to 1 : (1 : (0 + 1)) \\
&\to 1 : (1 : 1)
\end{aligned}
$$

Definition

Leftmost innermost strategy always reduces the leftmost of the innermost redexes.

Example (cont'd)

$$
\begin{aligned}
(0 : (1 + 2)) + (3 + 4) + (5 + 6) &\to ((0 : 3) + (3 + 4)) + (5 + 6) \\
&\to (3 + (3 + 4)) + (5 + 6) \\
&\to (3 + 7) + (5 + 6) \\
&\to (1 : 0) + (5 + 6) \\
&\to (1 : 0) + (1 : 1) \\
&\to 1 : (0 + (1 : 1)) \\
&\to 1 : (1 : (0 + 1)) \\
&\to 1 : (1 : 1) \\
&\to (1 + 1) : 1
\end{aligned}
$$

Definition

Leftmost innermost strategy always reduces the leftmost of the innermost redexes.

Example (cont'd)

$$
\begin{aligned}
(0 : (1 + 2)) + (3 + 4) + (5 + 6) &\to ((0 : 3) + (3 + 4)) + (5 + 6) \\
&\to (3 + (3 + 4)) + (5 + 6) \\
&\to (3 + 7) + (5 + 6) \\
&\to (1 : 0) + (5 + 6) \\
&\to (1 : 0) + (1 : 1) \\
&\to 1 : (0 + (1 : 1)) \\
&\to 1 : (1 : (0 + 1)) \\
&\to 1 : (1 : 1) \\
&\to (1 + 1) : 1 \\
&\to 2 : 1
\end{aligned}
$$

Definition

Parallel outermost strategy always reduces all outermost redexes in parallel.

Definition

Parallel outermost strategy always reduces all outermost redexes in parallel.

Example (cont'd)

$$((0 : (1 + 2)) + (3 + 4)) + (5 + 6)$$

Definition

Parallel outermost strategy always reduces all outermost redexes in parallel.

Example (cont'd)

$$((0 : (1 + 2)) + (3 + 4)) + (5 + 6) \Rightarrow\!\!\!\!\!\!+\;\; (0 : ((1 + 2) + (3 + 4))) + (1 : 1)$$

Definition

Parallel outermost strategy always reduces all outermost redexes in parallel.

Example (cont'd)

$$((0 : (1 + 2)) + (3 + 4)) + (5 + 6) \Rightarrow\!\!\!\!\!\!+\;\; (0 : ((1 + 2) + (3 + 4))) + (1 : 1)$$
$$\rightarrow 0 : (((1 + 2) + (3 + 4)) + (1 : 1))$$

Definition

Parallel outermost strategy always reduces all outermost redexes in parallel.

Example (cont'd)

$$((0 : (1 + 2)) + (3 + 4)) + (5 + 6) \Rightarrow\!\!\!\!\!\!+\ (0 : ((1 + 2) + (3 + 4))) + (1 : 1)$$
$$\rightarrow 0 : (((1 + 2) + (3 + 4)) + (1 : 1))$$
$$\rightarrow ((1 + 2) + (3 + 4)) + (1 : 1)$$

Definition

Parallel outermost strategy always reduces all outermost redexes in parallel.

Example (cont'd)

$$((0 : (1 + 2)) + (3 + 4)) + (5 + 6) \Rightarrow\!\!\!\!\!\!+\;\; (0 : ((1 + 2) + (3 + 4))) + (1 : 1)$$
$$\rightarrow 0 : (((1 + 2) + (3 + 4)) + (1 : 1))$$
$$\rightarrow ((1 + 2) + (3 + 4)) + (1 : 1)$$
$$\rightarrow 1 : (((1 + 2) + (3 + 4)) + 1)$$

Definition

Parallel outermost strategy always reduces all outermost redexes in parallel.

Example (cont'd)

$$
\begin{aligned}
((0 : (1 + 2)) + (3 + 4)) + (5 + 6) &\Mapsto (0 : ((1 + 2) + (3 + 4))) + (1 : 1) \\
&\to 0 : (((1 + 2) + (3 + 4)) + (1 : 1)) \\
&\to ((1 + 2) + (3 + 4)) + (1 : 1) \\
&\to 1 : (((1 + 2) + (3 + 4)) + 1) \\
&\Mapsto 1 : ((3 + 7) + 1)
\end{aligned}
$$

Definition

Parallel outermost strategy always reduces all outermost redexes in parallel.

Example (cont'd)

$$
\begin{aligned}
((0 : (1 + 2)) + (3 + 4)) + (5 + 6) &\Rightarrow\!\!\!\!\!\!\Rightarrow (0 : ((1 + 2) + (3 + 4))) + (1 : 1) \\
&\rightarrow 0 : (((1 + 2) + (3 + 4)) + (1 : 1)) \\
&\rightarrow ((1 + 2) + (3 + 4)) + (1 : 1) \\
&\rightarrow 1 : (((1 + 2) + (3 + 4)) + 1) \\
&\Rightarrow\!\!\!\!\!\!\Rightarrow 1 : ((3 + 7) + 1) \\
&\rightarrow 1 : ((1 : 0) + 1)
\end{aligned}
$$

Definition

Parallel outermost strategy always reduces all outermost redexes in parallel.

Example (cont'd)

$$
\begin{aligned}
((0 : (1 + 2)) + (3 + 4)) + (5 + 6) &\Mapsto (0 : ((1 + 2) + (3 + 4))) + (1 : 1) \\
&\to 0 : (((1 + 2) + (3 + 4)) + (1 : 1)) \\
&\to ((1 + 2) + (3 + 4)) + (1 : 1) \\
&\to 1 : (((1 + 2) + (3 + 4)) + 1) \\
&\Mapsto 1 : ((3 + 7) + 1) \\
&\to 1 : ((1 : 0) + 1) \\
&\to 1 : (1 : (0 + 1))
\end{aligned}
$$

Definition

Parallel outermost strategy always reduces all outermost redexes in parallel.

Example (cont'd)

$$
\begin{aligned}
((0 : (1 + 2)) + (3 + 4)) + (5 + 6) &\Rightarrow\!\!\!\!\!\!\Rightarrow (0 : ((1 + 2) + (3 + 4))) + (1 : 1) \\
&\rightarrow 0 : (((1 + 2) + (3 + 4)) + (1 : 1)) \\
&\rightarrow ((1 + 2) + (3 + 4)) + (1 : 1) \\
&\rightarrow 1 : (((1 + 2) + (3 + 4)) + 1) \\
&\Rightarrow\!\!\!\!\!\!\Rightarrow 1 : ((3 + 7) + 1) \\
&\rightarrow 1 : ((1 : 0) + 1) \\
&\rightarrow 1 : (1 : (0 + 1)) \\
&\rightarrow (1 + 1) : (0 + 1)
\end{aligned}
$$

Definition

Parallel outermost strategy always reduces all outermost redexes in parallel.

Example (cont'd)

$$
\begin{aligned}
((0 : (1 + 2)) + (3 + 4)) + (5 + 6) &\Rightarrow\!\!\!\!\!\!\Rightarrow (0 : ((1 + 2) + (3 + 4))) + (1 : 1) \\
&\rightarrow 0 : (((1 + 2) + (3 + 4)) + (1 : 1)) \\
&\rightarrow ((1 + 2) + (3 + 4)) + (1 : 1) \\
&\rightarrow 1 : (((1 + 2) + (3 + 4)) + 1) \\
&\Rightarrow\!\!\!\!\!\!\Rightarrow 1 : ((3 + 7) + 1) \\
&\rightarrow 1 : ((1 : 0) + 1) \\
&\rightarrow 1 : (1 : (0 + 1)) \\
&\rightarrow (1 + 1) : (0 + 1) \\
&\Rightarrow\!\!\!\!\!\!\Rightarrow 2 : 1
\end{aligned}
$$

Definition

Parallel innermost strategy always reduces all innermost redexes in parallel.

Definition

Parallel innermost strategy always reduces all innermost redexes in parallel.

Example (cont'd)

$$((0 : (1 + 2)) + (3 + 4)) + (5 + 6)$$

Definition

Parallel innermost strategy always reduces all innermost redexes in parallel.

Example (cont'd)

$$((0 : (1 + 2)) + (3 + 4)) + (5 + 6) \twoheadrightarrow ((0 : 3) + 7) + (1 : 1)$$

Definition

Parallel innermost strategy always reduces all innermost redexes in parallel.

Example (cont'd)

$$((0 : (1 + 2)) + (3 + 4)) + (5 + 6) \Rightarrow ((0 : 3) + 7) + (1 : 1)$$
$$\rightarrow (3 + 7) + (1 : 1)$$

Definition

Parallel innermost strategy always reduces all innermost redexes in parallel.

Example (cont'd)

$$((0 : (1 + 2)) + (3 + 4)) + (5 + 6) \Rightarrow ((0 : 3) + 7) + (1 : 1)$$
$$\rightarrow (3 + 7) + (1 : 1)$$
$$\rightarrow (1 : 0) + (1 : 1)$$

Definition

Parallel innermost strategy always reduces all innermost redexes in parallel.

Example (cont'd)

$$((0 : (1 + 2)) + (3 + 4)) + (5 + 6) \Rightarrow ((0 : 3) + 7) + (1 : 1)$$
$$\rightarrow (3 + 7) + (1 : 1)$$
$$\rightarrow (1 : 0) + (1 : 1)$$
$$\rightarrow 1 : (0 + (1 : 1))$$

Definition

Parallel innermost strategy always reduces all innermost redexes in parallel.

Example (cont'd)

$$((0 : (1 + 2)) + (3 + 4)) + (5 + 6) \Rightarrow ((0 : 3) + 7) + (1 : 1)$$
$$\rightarrow (3 + 7) + (1 : 1)$$
$$\rightarrow (1 : 0) + (1 : 1)$$
$$\rightarrow 1 : (0 + (1 : 1))$$
$$\rightarrow 1 : (1 : (0 + 1))$$

Definition

Parallel innermost strategy always reduces all innermost redexes in parallel.

Example (cont'd)

$$
\begin{aligned}
((0 : (1 + 2)) + (3 + 4)) + (5 + 6) &\Rightarrow\!\!\!\!\!\!\Rightarrow ((0 : 3) + 7) + (1 : 1) \\
&\rightarrow (3 + 7) + (1 : 1) \\
&\rightarrow (1 : 0) + (1 : 1) \\
&\rightarrow 1 : (0 + (1 : 1)) \\
&\rightarrow 1 : (1 : (0 + 1)) \\
&\rightarrow 1 : (1 : 1)
\end{aligned}
$$

Definition

Parallel innermost strategy always reduces all innermost redexes in parallel.

Example (cont'd)

$$
\begin{aligned}
((0 : (1 + 2)) + (3 + 4)) + (5 + 6) &\Rightarrow\!\!\!\!\!\Rightarrow ((0 : 3) + 7) + (1 : 1) \\
&\to (3 + 7) + (1 : 1) \\
&\to (1 : 0) + (1 : 1) \\
&\to 1 : (0 + (1 : 1)) \\
&\to 1 : (1 : (0 + 1)) \\
&\to 1 : (1 : 1) \\
&\to (1 + 1) : 1
\end{aligned}
$$

Definition

Parallel innermost strategy always reduces all innermost redexes in parallel.

Example (cont'd)

$$
\begin{aligned}
((0 : (1 + 2)) + (3 + 4)) + (5 + 6) &\Rightarrow\!\!\!\!\!\!\Rightarrow ((0 : 3) + 7) + (1 : 1) \\
&\rightarrow (3 + 7) + (1 : 1) \\
&\rightarrow (1 : 0) + (1 : 1) \\
&\rightarrow 1 : (0 + (1 : 1)) \\
&\rightarrow 1 : (1 : (0 + 1)) \\
&\rightarrow 1 : (1 : 1) \\
&\rightarrow (1 + 1) : 1 \\
&\rightarrow 2 : 1
\end{aligned}
$$

Definition

A development of set of redex positions Q in term t is a rewrite sequence starting from t in which all contracted redex positions descend from positions in Q.

Example

- rewrite rules

$$0 + y \to y \qquad 0 \times y \to 0$$
$$\mathsf{s}(x) + y \to \mathsf{s}(x + y) \qquad \mathsf{s}(x) \times y \to (x \times y) + y$$

- rewrite sequences

$$\underline{\mathsf{s}(0) \times (0 \times 0)} \to (0 \times \underline{(0 \times 0)}) + (0 \times 0) \to (0 \times 0) + (0 \times 0)$$ ☺

Definition

A development of set of redex positions Q in term t is a rewrite sequence starting from t in which all contracted redex positions descend from positions in Q.

Example

- rewrite rules

$$0 + y \to y \qquad 0 \times y \to 0$$
$$s(x) + y \to s(x + y) \qquad s(x) \times y \to (x \times y) + y$$

- rewrite sequences

$$\underline{s(0) \times (0 \times 0)} \to (0 \times \underline{(0 \times 0)}) + (0 \times 0) \to (0 \times 0) + (0 \times 0) \quad ☺$$
$$\underline{s(0) \times (0 \times 0)} \to \underline{(0 \times (0 \times 0))} + (0 \times 0) \to 0 + (0 \times 0) \quad ☹$$

Definition

A development of set of redex positions Q in term t is a rewrite sequence starting from t in which all contracted redex positions descend from positions in Q.

Example

- rewrite rules

$$0 + y \to y \qquad 0 \times y \to 0$$
$$\mathsf{s}(x) + y \to \mathsf{s}(x + y) \qquad \mathsf{s}(x) \times y \to (x \times y) + y$$

- rewrite sequences

$$\underline{\mathsf{s}(0) \times (0 \times 0)} \to (0 \times \underline{(0 \times 0)}) + (0 \times 0) \to (0 \times 0) + (0 \times 0) \quad ☺$$
$$\underline{\mathsf{s}(0) \times (0 \times 0)} \to \underline{(0 \times (0 \times 0))} + (0 \times 0) \to 0 + (0 \times 0) \quad ☹$$
$$\mathsf{s}(0) \times \underline{(0 \times 0)} \to \underline{\mathsf{s}(0) \times 0} \to (0 \times 0) + 0 \quad ☺$$

Definition (Overlining)

For a TRS $\mathcal{R} = \langle \Sigma, R \rangle$ we define the overlined TRS $\overline{\mathcal{R}} = \langle \overline{\Sigma}, \overline{R} \rangle$:

- $\overline{\Sigma} = \Sigma \cup \{\overline{f} \mid f \in \Sigma\}$,
- $\overline{R} = \{\overline{\rho} \mid \rho \in R\}$

where $\overline{\rho}$ is obtained from ρ by overlining the head symbol of the left-hand side.

$$\rho : f(s_1, \ldots, s_n) \to r \quad \text{yields} \quad \overline{\rho} : \overline{f}(s_1, \ldots, s_n) \to r$$

Definition (Overlining)

For a TRS $\mathcal{R} = \langle \Sigma, R \rangle$ we define the overlined TRS $\overline{\mathcal{R}} = \langle \overline{\Sigma}, \overline{R} \rangle$:

- $\overline{\Sigma} = \Sigma \cup \{\overline{f} \mid f \in \Sigma\}$,
- $\overline{R} = \{\overline{\rho} \mid \rho \in R\}$

where $\overline{\rho}$ is obtained from ρ by overlining the head symbol of the left-hand side.

$$\rho : f(s_1, \ldots, s_n) \to r \quad \text{yields} \quad \overline{\rho} : \overline{f}(s_1, \ldots, s_n) \to r$$

Example

The overlined version of Combinatory Logic:

$$\begin{array}{lcl} \overline{Ap}(Ap(Ap(S,x),y),z) & \to & Ap(Ap(x,z),Ap(y,z)) \\ \overline{Ap}(Ap(K,x),y) & \to & x \\ \overline{Ap}(I,x) & \to & x \end{array}$$

We write $t \geq s$ if t can be obtained from s by overlining some redex positions.

Definition (Lifting)

A $\mathcal{R}$ rewrite sequence $A : s_1 \to s_2 \to \ldots \to s_n$ can be lifted if:

$$\begin{array}{ccccccc}
t_1 & \xrightarrow{\langle \overline{\rho}_1,\, p_1 \rangle} & t_2 & \xrightarrow{\langle \overline{\rho}_2,\, p_2 \rangle} & \cdots & \xrightarrow{\langle \overline{\rho}_{n-1},\, p_{n-1} \rangle} & t_n \\
\vee| & & \vee| & & & & \vee| \\
s_1 & \xrightarrow{\langle \rho_1,\, p_1 \rangle} & s_2 & \xrightarrow{\langle \rho_2,\, p_2 \rangle} & \cdots & \xrightarrow{\langle \rho_{n-1},\, p_{n-1} \rangle} & s_n
\end{array}$$

for some $\overline{\mathcal{R}}$ rewrite sequence $B : t_1 \to \ldots t_n$.

We write $t \geq s$ if t can be obtained from s by overlining some redex positions.

Definition (Lifting)

A $\mathcal{R}$ rewrite sequence $A : s_1 \to s_2 \to \ldots \to s_n$ can be lifted if:

$$\begin{array}{ccccccc}
t_1 & \xrightarrow{\langle \overline{\rho}_1,\, p_1 \rangle} & t_2 & \xrightarrow{\langle \overline{\rho}_2,\, p_2 \rangle} & \cdots & \xrightarrow{\langle \overline{\rho}_{n-1},\, p_{n-1} \rangle} & t_n \\
\vee| & & \vee| & & & & \vee| \\
s_1 & \xrightarrow{\langle \rho_1,\, p_1 \rangle} & s_2 & \xrightarrow{\langle \rho_2,\, p_2 \rangle} & \cdots & \xrightarrow{\langle \rho_{n-1},\, p_{n-1} \rangle} & s_n
\end{array}$$

for some $\overline{\mathcal{R}}$ rewrite sequence $B : t_1 \to \ldots t_n$.

Lemma

For orthogonal TRSs: a reduction is a development $\iff$ *it can be lifted.*

Theorem

Properties of $\overline{\mathcal{R}}$ for orthogonal TRSs $\mathcal{R}$:

- *$\overline{\mathcal{R}}$ is orthogonal.*
- *$\overline{\mathcal{R}}$ is SN.*
- *$\overline{\mathcal{R}}$ is CR.*

For SN we show that $\overline{\mathcal{R}}$ is ILPO terminating where $\overline{f} > g$ for every $f, g \in \Sigma$.

Let $\ell = \overline{f}(\ell_1, \ldots, \ell_n)$ and $r \in \mathcal{T}(\Sigma, \mathcal{X})$ with $\mathcal{V}\mathrm{ar}(r) \subseteq \mathcal{V}\mathrm{ar}\ell$.

Then $\ell^* \to^*_{ilpo} r$ by induction on r:

- If $r \in \mathcal{V}\mathrm{ar}(\ell)$, then we use $\to_{put}$ and $\to_{select}$.
- If $r = g(r_1, \ldots, r_m)$, then we use $\ell^* \to_{copy} g(\ell^*, \ldots, \ell^*)$.
  Moreover by the induction hypothesis $\ell^* \to^*_{ilpo} r_i$ for every i.

Theorem

Properties of $\overline{\mathcal{R}}$ for orthogonal TRSs $\mathcal{R}$:

- *$\overline{\mathcal{R}}$ is orthogonal.*
- *$\overline{\mathcal{R}}$ is SN.*
- *$\overline{\mathcal{R}}$ is CR.*

Proof.

The orthogonality of $\overline{\mathcal{R}}$ is immediate. Hence $\overline{\mathcal{R}}$ is CR.

Theorem

Properties of $\overline{\mathcal{R}}$ for orthogonal TRSs $\mathcal{R}$:

- *$\overline{\mathcal{R}}$ is orthogonal.*
- *$\overline{\mathcal{R}}$ is SN.*
- *$\overline{\mathcal{R}}$ is CR.*

Proof.

The orthogonality of $\overline{\mathcal{R}}$ is immediate. Hence $\overline{\mathcal{R}}$ is CR.

For SN we show that $\overline{\mathcal{R}}$ is ILPO terminating where $\overline{f} > g$ for every $f, g \in \Sigma$.

Theorem

Properties of $\overline{\mathcal{R}}$ for orthogonal TRSs $\mathcal{R}$:

- *$\overline{\mathcal{R}}$ is orthogonal.*
- *$\overline{\mathcal{R}}$ is SN.*
- *$\overline{\mathcal{R}}$ is CR.*

Proof.

The orthogonality of $\overline{\mathcal{R}}$ is immediate. Hence $\overline{\mathcal{R}}$ is CR.

For SN we show that $\overline{\mathcal{R}}$ is ILPO terminating where $\overline{f} > g$ for every $f, g \in \Sigma$.
Let $\ell = \overline{f}(\ell_1, \ldots, \ell_n)$ and $r \in \mathcal{T}(\Sigma, \mathcal{X})$ with $\mathcal{V}\mathrm{ar}(r) \subseteq \mathcal{V}\mathrm{ar}\ell$.

Theorem

Properties of $\overline{\mathcal{R}}$ for orthogonal TRSs $\mathcal{R}$:

- *$\overline{\mathcal{R}}$ is orthogonal.*
- *$\overline{\mathcal{R}}$ is SN.*
- *$\overline{\mathcal{R}}$ is CR.*

Proof.

The orthogonality of $\overline{\mathcal{R}}$ is immediate. Hence $\overline{\mathcal{R}}$ is CR.

For SN we show that $\overline{\mathcal{R}}$ is ILPO terminating where $\overline{f} > g$ for every $f, g \in \Sigma$.
Let $\ell = \overline{f}(\ell_1, \ldots, \ell_n)$ and $r \in \mathcal{T}(\Sigma, \mathcal{X})$ with $\mathcal{V}\mathrm{ar}(r) \subseteq \mathcal{V}\mathrm{ar}\ell$.
Then $\ell^* \to^*_{ilpo} r$ by induction on r:

Theorem

Properties of $\overline{\mathcal{R}}$ for orthogonal TRSs $\mathcal{R}$:

- *$\overline{\mathcal{R}}$ is orthogonal.*
- *$\overline{\mathcal{R}}$ is SN.*
- *$\overline{\mathcal{R}}$ is CR.*

Proof.

The orthogonality of $\overline{\mathcal{R}}$ is immediate. Hence $\overline{\mathcal{R}}$ is CR.

For SN we show that $\overline{\mathcal{R}}$ is ILPO terminating where $\overline{f} > g$ for every $f, g \in \Sigma$.
Let $\ell = \overline{f}(\ell_1, \ldots, \ell_n)$ and $r \in \mathcal{T}(\Sigma, \mathcal{X})$ with $\mathcal{V}\mathrm{ar}(r) \subseteq \mathcal{V}\mathrm{ar}\ell$.
Then $\ell^* \to^*_{ilpo} r$ by induction on r:

- If $r \in \mathcal{V}\mathrm{ar}(\ell)$, then we use $\to_{put}$ and $\to_{select}$.

Theorem

Properties of $\overline{\mathcal{R}}$ for orthogonal TRSs $\mathcal{R}$:

- *$\overline{\mathcal{R}}$ is orthogonal.*
- *$\overline{\mathcal{R}}$ is SN.*
- *$\overline{\mathcal{R}}$ is CR.*

Proof.

The orthogonality of $\overline{\mathcal{R}}$ is immediate. Hence $\overline{\mathcal{R}}$ is CR.

For SN we show that $\overline{\mathcal{R}}$ is ILPO terminating where $\overline{f} > g$ for every $f, g \in \Sigma$.
Let $\ell = \overline{f}(\ell_1, \ldots, \ell_n)$ and $r \in \mathcal{T}(\Sigma, \mathcal{X})$ with $\mathcal{V}\mathrm{ar}(r) \subseteq \mathcal{V}\mathrm{ar}\ell$.
Then $\ell^* \to^*_{ilpo} r$ by induction on r:

- If $r \in \mathcal{V}\mathrm{ar}(\ell)$, then we use $\to_{put}$ and $\to_{select}$.
- If $r = g(r_1, \ldots, r_m)$, then we use $\ell^* \to_{copy} g(\ell^*, \ldots, \ell^*)$.

Theorem

Properties of $\overline{\mathcal{R}}$ for orthogonal TRSs $\mathcal{R}$:

- *$\overline{\mathcal{R}}$ is orthogonal.*
- *$\overline{\mathcal{R}}$ is SN.*
- *$\overline{\mathcal{R}}$ is CR.*

Proof.

The orthogonality of $\overline{\mathcal{R}}$ is immediate. Hence $\overline{\mathcal{R}}$ is CR.

For SN we show that $\overline{\mathcal{R}}$ is ILPO terminating where $\overline{f} > g$ for every $f, g \in \Sigma$.
Let $\ell = \overline{f}(\ell_1, \ldots, \ell_n)$ and $r \in \mathcal{T}(\Sigma, \mathcal{X})$ with $\mathcal{V}\mathrm{ar}(r) \subseteq \mathcal{V}\mathrm{ar}\ell$.
Then $\ell^* \to^*_{ilpo} r$ by induction on r:

- If $r \in \mathcal{V}\mathrm{ar}(\ell)$, then we use $\to_{put}$ and $\to_{select}$.
- If $r = g(r_1, \ldots, r_m)$, then we use $\ell^* \to_{copy} g(\ell^*, \ldots, \ell^*)$.
  Moreover by the induction hypothesis $\ell^* \to^*_{ilpo} r_i$ for every i.

Theorem

Developments are finite.

Theorem

Developments are finite.

Definition

A development $A\colon s \to^* t$ of $Q \subseteq \mathcal{P}\mathrm{os}(s)$ is complete if $Q/A = \varnothing$.

We write $s \mathrel{-\!\!\circ\!\!\to} t$ (called multi-step) if there is a complete development $s \to^* t$.

Theorem

Developments are finite.

Definition

A development $A\colon s \to^* t$ of $Q \subseteq \mathcal{P}\mathrm{os}(s)$ is complete if $Q/A = \varnothing$.

We write $s \mathrel{-\!\!\circ\!\!\to} t$ (called multi-step) if there is a complete development $s \to^* t$.

Example

$$\underline{\mathsf{s}(0) \times (0 \times 0)} \to (0 \times \underline{(0 \times 0)}) + (0 \times 0) \to (0 \times 0) + (0 \times 0) \qquad \text{☹}$$

$$\mathsf{s}(0) \times \underline{(0 \times 0)} \to \underline{\mathsf{s}(0) \times 0} \to (0 \times 0) + 0 \qquad \text{☺}$$

Theorem

Developments are finite.

Definition

A development $A\colon s \to^* t$ of $Q \subseteq \mathcal{P}\mathrm{os}(s)$ is complete if $Q/A = \varnothing$.

We write $s \twoheadrightarrow t$ (called multi-step) if there is a complete development $s \to^* t$.

Example

$$\underline{\mathsf{s}(0) \times (0 \times 0)} \to (0 \times \underline{(0 \times 0)}) + (0 \times 0) \to (0 \times 0) + (0 \times 0) \qquad \text{☹}$$

$$\mathsf{s}(0) \times \underline{(0 \times 0)} \to \underline{\mathsf{s}(0) \times 0} \to (0 \times 0) + 0 \qquad \text{☺}$$

Theorem

All complete developments of Q are permutation equivalent.

Definition

For orthogonal TRSs the full substitution strategy performs complete development of all redexes.

Definition

For orthogonal TRSs the full substitution strategy performs complete development of all redexes.

Example

- rewrite rules

$$0 + y \to y \qquad 0 \times y \to 0$$
$$\mathsf{s}(x) + y \to \mathsf{s}(x + y) \qquad \mathsf{s}(x) \times y \to (x \times y) + y$$

- full substitution strategy

$$\mathsf{s}(\mathsf{s}(0)) \times (\mathsf{s}(0) + \mathsf{s}(\mathsf{s}(0)))$$

Definition

For orthogonal TRSs the full substitution strategy performs complete development of all redexes.

Example

- rewrite rules

$$0 + y \to y \qquad 0 \times y \to 0$$
$$\mathsf{s}(x) + y \to \mathsf{s}(x + y) \qquad \mathsf{s}(x) \times y \to (x \times y) + y$$

- full substitution strategy

$$\mathsf{s}(\mathsf{s}(0)) \times (\mathsf{s}(0) + \mathsf{s}(\mathsf{s}(0)))$$
$$\twoheadrightarrow (\mathsf{s}(0) \times \mathsf{s}(0 + \mathsf{s}(\mathsf{s}(0)))) + \mathsf{s}(0 + \mathsf{s}(\mathsf{s}(0)))$$

Definition

For orthogonal TRSs the full substitution strategy performs complete development of all redexes.

Example

- rewrite rules

$$0 + y \to y \qquad 0 \times y \to 0$$
$$\mathsf{s}(x) + y \to \mathsf{s}(x + y) \qquad \mathsf{s}(x) \times y \to (x \times y) + y$$

- full substitution strategy

$$\begin{aligned} &\mathsf{s}(\mathsf{s}(0)) \times (\mathsf{s}(0) + \mathsf{s}(\mathsf{s}(0))) \\ &\quad \twoheadrightarrow (\mathsf{s}(0) \times \mathsf{s}(0 + \mathsf{s}(\mathsf{s}(0)))) + \mathsf{s}(0 + \mathsf{s}(\mathsf{s}(0))) \\ &\quad \twoheadrightarrow ((0 \times \mathsf{s}(\mathsf{s}(\mathsf{s}(0)))) + \mathsf{s}(\mathsf{s}(\mathsf{s}(0)))) + \mathsf{s}(\mathsf{s}(0))) \end{aligned}$$

Definition

For orthogonal TRSs the full substitution strategy performs complete development of all redexes.

Example

- rewrite rules

$$0 + y \to y \qquad 0 \times y \to 0$$
$$\mathsf{s}(x) + y \to \mathsf{s}(x + y) \qquad \mathsf{s}(x) \times y \to (x \times y) + y$$

- full substitution strategy

$$\begin{aligned}
&\mathsf{s}(\mathsf{s}(0)) \times (\mathsf{s}(0) + \mathsf{s}(\mathsf{s}(0))) \\
&\quad \twoheadrightarrow (\mathsf{s}(0) \times \mathsf{s}(0 + \mathsf{s}(\mathsf{s}(0)))) + \mathsf{s}(0 + \mathsf{s}(\mathsf{s}(0))) \\
&\quad \twoheadrightarrow ((0 \times \mathsf{s}(\mathsf{s}(\mathsf{s}(0)))) + \mathsf{s}(\mathsf{s}(\mathsf{s}(0)))) + \mathsf{s}(\mathsf{s}(\mathsf{s}(0))) \\
&\quad \to (0 + \mathsf{s}(\mathsf{s}(\mathsf{s}(0)))) + \mathsf{s}(\mathsf{s}(\mathsf{s}(0)))
\end{aligned}$$

Definition

For orthogonal TRSs the full substitution strategy performs complete development of all redexes.

Example

- rewrite rules

$$0 + y \to y \qquad 0 \times y \to 0$$
$$\mathsf{s}(x) + y \to \mathsf{s}(x + y) \qquad \mathsf{s}(x) \times y \to (x \times y) + y$$

- full substitution strategy

$$\begin{aligned}
&\mathsf{s}(\mathsf{s}(0)) \times (\mathsf{s}(0) + \mathsf{s}(\mathsf{s}(0))) \\
&\quad \twoheadrightarrow (\mathsf{s}(0) \times \mathsf{s}(0 + \mathsf{s}(\mathsf{s}(0)))) + \mathsf{s}(0 + \mathsf{s}(\mathsf{s}(0))) \\
&\quad \twoheadrightarrow ((0 \times \mathsf{s}(\mathsf{s}(\mathsf{s}(0)))) + \mathsf{s}(\mathsf{s}(\mathsf{s}(0)))) + \mathsf{s}(\mathsf{s}(0))) \\
&\quad \to (0 + \mathsf{s}(\mathsf{s}(\mathsf{s}(0)))) + \mathsf{s}(\mathsf{s}(0))) \\
&\quad \to \mathsf{s}(\mathsf{s}(\mathsf{s}(0))) + \mathsf{s}(\mathsf{s}(0)))
\end{aligned}$$

Definition

For orthogonal TRSs the full substitution strategy performs complete development of all redexes.

Example

- rewrite rules

$$0 + y \to y \qquad 0 \times y \to 0$$
$$\mathsf{s}(x) + y \to \mathsf{s}(x + y) \qquad \mathsf{s}(x) \times y \to (x \times y) + y$$

- full substitution strategy

$$\begin{aligned}
&\mathsf{s}(\mathsf{s}(0)) \times (\mathsf{s}(0) + \mathsf{s}(\mathsf{s}(0))) \\
&\quad \twoheadrightarrow (\mathsf{s}(0) \times \mathsf{s}(0 + \mathsf{s}(\mathsf{s}(0)))) + \mathsf{s}(0 + \mathsf{s}(\mathsf{s}(0))) \\
&\quad \twoheadrightarrow ((0 \times \mathsf{s}(\mathsf{s}(\mathsf{s}(0)))) + \mathsf{s}(\mathsf{s}(\mathsf{s}(0)))) + \mathsf{s}(\mathsf{s}(0))) \\
&\quad \to (0 + \mathsf{s}(\mathsf{s}(\mathsf{s}(0)))) + \mathsf{s}(\mathsf{s}(0))) \\
&\quad \to \mathsf{s}(\mathsf{s}(\mathsf{s}(0))) + \mathsf{s}(\mathsf{s}(0))) \\
&\quad \to \ldots \to \mathsf{s}(\mathsf{s}(\mathsf{s}(\mathsf{s}(\mathsf{s}(0))))))
\end{aligned}$$

Outline

- Overview
- Strategies
  - Definitions
  - Results

Theorem

For orthogonal TRSs

- *full substitution* and *parallel outermost* strategies are *normalizing*

Theorem

For orthogonal TRSs

- *full substitution* and *parallel outermost* strategies are *normalizing*
- *innermost* strategies are *perpetual*

Theorem

For orthogonal TRSs

- *full substitution* and *parallel outermost* strategies are *normalizing*
- *innermost* strategies are *perpetual*
- *leftmost outermost* strategy is *not normalizing*

Theorem

For orthogonal TRSs

- *full substitution* and *parallel outermost* strategies are *normalizing*
- *innermost* strategies are *perpetual*
- *leftmost outermost* strategy is *not normalizing*
- *full substitution* is *fair*

Example

$$\mathsf{a} \to \mathsf{b} \qquad \mathsf{c} \to \mathsf{c} \qquad \mathsf{f}(x, \mathsf{b}) \to \mathsf{b}$$

- leftmost outermost $\mathsf{f}(\mathsf{c}, \mathsf{a})$
- leftmost innermost $\mathsf{f}(\mathsf{c}, \mathsf{a})$
- parallel outermost $\mathsf{f}(\mathsf{c}, \mathsf{a})$
- parallel innermost $\mathsf{f}(\mathsf{c}, \mathsf{a})$
- full substitution $\mathsf{f}(\mathsf{c}, \mathsf{a})$

Theorem

For orthogonal TRSs

- *full substitution* and *parallel outermost* strategies are *normalizing*
- *innermost* strategies are *perpetual*
- *leftmost outermost* strategy is *not normalizing*
- *full substitution* is *fair*

Example

$a \to b$ $\quad c \to c$ $\quad f(x, b) \to b$

- leftmost outermost $\quad f(c, a) \to f(c, a)$
- leftmost innermost $\quad f(c, a)$
- parallel outermost $\quad f(c, a)$
- parallel innermost $\quad f(c, a)$
- full substitution $\quad f(c, a)$

Theorem

For orthogonal TRSs

- *full substitution* and *parallel outermost* strategies are *normalizing*
- *innermost* strategies are *perpetual*
- *leftmost outermost* strategy is *not normalizing*
- *full substitution* is *fair*

Example

$$a \to b \qquad c \to c \qquad f(x, b) \to b$$

- leftmost outermost $f(c, a) \to f(c, a) \to f(c, a)$
- leftmost innermost $f(c, a)$
- parallel outermost $f(c, a)$
- parallel innermost $f(c, a)$
- full substitution $f(c, a)$

Theorem

For orthogonal TRSs

- *full substitution* and *parallel outermost* strategies are *normalizing*
- *innermost* strategies are *perpetual*
- *leftmost outermost* strategy is *not normalizing*
- *full substitution* is *fair*

Example

$$\mathsf{a} \to \mathsf{b} \qquad \mathsf{c} \to \mathsf{c} \qquad \mathsf{f}(x, \mathsf{b}) \to \mathsf{b}$$

- leftmost outermost $\mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \cdots$
- leftmost innermost $\mathsf{f}(\mathsf{c}, \mathsf{a})$
- parallel outermost $\mathsf{f}(\mathsf{c}, \mathsf{a})$
- parallel innermost $\mathsf{f}(\mathsf{c}, \mathsf{a})$
- full substitution $\mathsf{f}(\mathsf{c}, \mathsf{a})$

Theorem

For orthogonal TRSs

- *full substitution* and *parallel outermost* strategies are *normalizing*
- *innermost* strategies are *perpetual*
- *leftmost outermost* strategy is *not normalizing*
- *full substitution* is *fair*

Example

$$\mathsf{a} \to \mathsf{b} \qquad \mathsf{c} \to \mathsf{c} \qquad \mathsf{f}(x, \mathsf{b}) \to \mathsf{b}$$

- leftmost outermost $\mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \cdots$
- leftmost innermost $\mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a})$
- parallel outermost $\mathsf{f}(\mathsf{c}, \mathsf{a})$
- parallel innermost $\mathsf{f}(\mathsf{c}, \mathsf{a})$
- full substitution $\mathsf{f}(\mathsf{c}, \mathsf{a})$

Theorem

For orthogonal TRSs

- *full substitution* and *parallel outermost* strategies are *normalizing*
- *innermost* strategies are *perpetual*
- *leftmost outermost* strategy is *not normalizing*
- *full substitution* is *fair*

Example

$$\mathsf{a} \to \mathsf{b} \qquad \mathsf{c} \to \mathsf{c} \qquad \mathsf{f}(x, \mathsf{b}) \to \mathsf{b}$$

- leftmost outermost $\mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \cdots$
- leftmost innermost $\mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a})$
- parallel outermost $\mathsf{f}(\mathsf{c}, \mathsf{a})$
- parallel innermost $\mathsf{f}(\mathsf{c}, \mathsf{a})$
- full substitution $\mathsf{f}(\mathsf{c}, \mathsf{a})$

Theorem

For orthogonal TRSs

- *full substitution* and *parallel outermost* strategies are *normalizing*
- *innermost* strategies are *perpetual*
- *leftmost outermost* strategy is *not normalizing*
- *full substitution* is *fair*

Example

$$\mathsf{a} \to \mathsf{b} \qquad \mathsf{c} \to \mathsf{c} \qquad \mathsf{f}(x, \mathsf{b}) \to \mathsf{b}$$

- leftmost outermost $\mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \cdots$
- leftmost innermost $\mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \cdots$
- parallel outermost $\mathsf{f}(\mathsf{c}, \mathsf{a})$
- parallel innermost $\mathsf{f}(\mathsf{c}, \mathsf{a})$
- full substitution $\mathsf{f}(\mathsf{c}, \mathsf{a})$

Theorem

For orthogonal TRSs

- *full substitution* and *parallel outermost* strategies are *normalizing*
- *innermost* strategies are *perpetual*
- *leftmost outermost* strategy is *not normalizing*
- *full substitution* is *fair*

Example

$$\mathsf{a} \to \mathsf{b} \qquad \mathsf{c} \to \mathsf{c} \qquad \mathsf{f}(x, \mathsf{b}) \to \mathsf{b}$$

- leftmost outermost $\mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \cdots$
- leftmost innermost $\mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \cdots$
- parallel outermost $\mathsf{f}(\mathsf{c}, \mathsf{a}) \mathrel{+\!\!\!\to} \mathsf{f}(\mathsf{c}, \mathsf{b})$
- parallel innermost $\mathsf{f}(\mathsf{c}, \mathsf{a})$
- full substitution $\mathsf{f}(\mathsf{c}, \mathsf{a})$

Theorem

For orthogonal TRSs

- *full substitution* and *parallel outermost* strategies are *normalizing*
- *innermost* strategies are *perpetual*
- *leftmost outermost* strategy is *not normalizing*
- *full substitution* is *fair*

Example

$$\mathsf{a} \to \mathsf{b} \qquad \mathsf{c} \to \mathsf{c} \qquad \mathsf{f}(x, \mathsf{b}) \to \mathsf{b}$$

- leftmost outermost $\quad \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \cdots$
- leftmost innermost $\quad \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \cdots$
- parallel outermost $\quad \mathsf{f}(\mathsf{c}, \mathsf{a}) \mathrel{+\!\!\!\to} \mathsf{f}(\mathsf{c}, \mathsf{b}) \to \mathsf{b}$
- parallel innermost $\quad \mathsf{f}(\mathsf{c}, \mathsf{a})$
- full substitution $\quad \mathsf{f}(\mathsf{c}, \mathsf{a})$

Theorem

For orthogonal TRSs

- *full substitution* and *parallel outermost* strategies are *normalizing*
- *innermost* strategies are *perpetual*
- *leftmost outermost* strategy is *not normalizing*
- *full substitution* is *fair*

Example

$$\mathsf{a} \to \mathsf{b} \qquad \mathsf{c} \to \mathsf{c} \qquad \mathsf{f}(x, \mathsf{b}) \to \mathsf{b}$$

- leftmost outermost $\mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \cdots$
- leftmost innermost $\mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \cdots$
- parallel outermost $\mathsf{f}(\mathsf{c}, \mathsf{a}) \Rightarrow \mathsf{f}(\mathsf{c}, \mathsf{b}) \to \mathsf{b}$
- parallel innermost $\mathsf{f}(\mathsf{c}, \mathsf{a}) \Rightarrow \mathsf{f}(\mathsf{c}, \mathsf{b})$
- full substitution $\mathsf{f}(\mathsf{c}, \mathsf{a})$

Theorem

For orthogonal TRSs

- *full substitution* and *parallel outermost* strategies are *normalizing*
- *innermost* strategies are *perpetual*
- *leftmost outermost* strategy is *not normalizing*
- *full substitution* is *fair*

Example

$$\mathsf{a} \to \mathsf{b} \qquad \mathsf{c} \to \mathsf{c} \qquad \mathsf{f}(x, \mathsf{b}) \to \mathsf{b}$$

- leftmost outermost $\quad \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \cdots$
- leftmost innermost $\quad \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \cdots$
- parallel outermost $\quad \mathsf{f}(\mathsf{c}, \mathsf{a}) \mathrel{+\!\!\!\to} \mathsf{f}(\mathsf{c}, \mathsf{b}) \to \mathsf{b}$
- parallel innermost $\quad \mathsf{f}(\mathsf{c}, \mathsf{a}) \mathrel{+\!\!\!\to} \mathsf{f}(\mathsf{c}, \mathsf{b}) \to \mathsf{f}(\mathsf{c}, \mathsf{b})$
- full substitution $\quad \mathsf{f}(\mathsf{c}, \mathsf{a})$

Theorem

For orthogonal TRSs

- *full substitution* and *parallel outermost* strategies are *normalizing*
- *innermost* strategies are *perpetual*
- *leftmost outermost* strategy is *not normalizing*
- *full substitution* is *fair*

Example

$$\mathsf{a} \to \mathsf{b} \qquad \mathsf{c} \to \mathsf{c} \qquad \mathsf{f}(x, \mathsf{b}) \to \mathsf{b}$$

- leftmost outermost $\quad \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \cdots$
- leftmost innermost $\quad \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \cdots$
- parallel outermost $\quad \mathsf{f}(\mathsf{c}, \mathsf{a}) \Rightarrow \mathsf{f}(\mathsf{c}, \mathsf{b}) \to \mathsf{b}$
- parallel innermost $\quad \mathsf{f}(\mathsf{c}, \mathsf{a}) \Rightarrow \mathsf{f}(\mathsf{c}, \mathsf{b}) \to \mathsf{f}(\mathsf{c}, \mathsf{b}) \to \cdots$
- full substitution $\quad \mathsf{f}(\mathsf{c}, \mathsf{a})$

Theorem

For orthogonal TRSs

- *full substitution* and *parallel outermost* strategies are *normalizing*
- *innermost* strategies are *perpetual*
- *leftmost outermost* strategy is *not normalizing*
- *full substitution* is *fair*

Example

$$\mathsf{a} \to \mathsf{b} \qquad \mathsf{c} \to \mathsf{c} \qquad \mathsf{f}(x, \mathsf{b}) \to \mathsf{b}$$

- leftmost outermost $\quad \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \cdots$
- leftmost innermost $\quad \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \cdots$
- parallel outermost $\quad \mathsf{f}(\mathsf{c}, \mathsf{a}) \Rightarrow \mathsf{f}(\mathsf{c}, \mathsf{b}) \to \mathsf{b}$
- parallel innermost $\quad \mathsf{f}(\mathsf{c}, \mathsf{a}) \Rightarrow \mathsf{f}(\mathsf{c}, \mathsf{b}) \to \mathsf{f}(\mathsf{c}, \mathsf{b}) \to \cdots$
- full substitution $\quad \mathsf{f}(\mathsf{c}, \mathsf{a}) \Rightarrow \mathsf{f}(\mathsf{c}, \mathsf{b})$

Theorem

For orthogonal TRSs

- *full substitution* and *parallel outermost* strategies are *normalizing*
- *innermost* strategies are *perpetual*
- *leftmost outermost* strategy is *not normalizing*
- *full substitution* is *fair*

Example

$$\mathsf{a} \to \mathsf{b} \qquad \mathsf{c} \to \mathsf{c} \qquad \mathsf{f}(x, \mathsf{b}) \to \mathsf{b}$$

- leftmost outermost $\quad \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \cdots$
- leftmost innermost $\quad \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \mathsf{f}(\mathsf{c}, \mathsf{a}) \to \cdots$
- parallel outermost $\quad \mathsf{f}(\mathsf{c}, \mathsf{a}) \Rightarrow \mathsf{f}(\mathsf{c}, \mathsf{b}) \to \mathsf{b}$
- parallel innermost $\quad \mathsf{f}(\mathsf{c}, \mathsf{a}) \Rightarrow \mathsf{f}(\mathsf{c}, \mathsf{b}) \to \mathsf{f}(\mathsf{c}, \mathsf{b}) \to \cdots$
- full substitution $\quad \mathsf{f}(\mathsf{c}, \mathsf{a}) \Rightarrow \mathsf{f}(\mathsf{c}, \mathsf{b}) \Rightarrow \mathsf{b}$

Definition

A reduction $\rho = t_0 \to t_1 \to \ldots$ is cofinal if for every $t_0 \to^* s$ there exists t_n in ρ such that $s \to^* t_n$.

Let t_0 be a term that has a normal form u ($t_0 \to^* u$).

Consider a maximal $\mathcal{S}$ rewrite sequence $\rho : t_0 \to t_1 \to \ldots$ starting from t_0.

By cofinality there must be t_n in ρ such that $u \to^* t_n$.

Hence $t_n = u$ since u is a normal form.

Definition

A reduction $\rho = t_0 \to t_1 \to \ldots$ is cofinal if for every $t_0 \to^* s$ there exists t_n in ρ such that $s \to^* t_n$.

Definition

A strategy $\mathcal{S}$ is cofinal if every every maximal $\mathcal{S}$ rewrite sequence is cofinal.

Let t_0 be a term that has a normal form u ($t_0 \to^* u$).

Consider a maximal $\mathcal{S}$ rewrite sequence $\rho : t_0 \to t_1 \to \ldots$ starting from t_0.

By cofinality there must be t_n in ρ such that $u \to^* t_n$.

Hence $t_n = u$ since u is a normal form.

Definition

A reduction $\rho = t_0 \to t_1 \to \ldots$ is cofinal if for every $t_0 \to^* s$ there exists t_n in ρ such that $s \to^* t_n$.

Definition

A strategy $\mathcal{S}$ is cofinal if every every maximal $\mathcal{S}$ rewrite sequence is cofinal.

Theorem

Cofinal strategies are normalizing.

Let t_0 be a term that has a normal form u ($t_0 \to^* u$).

Consider a maximal $\mathcal{S}$ rewrite sequence $\rho : t_0 \to t_1 \to \ldots$ starting from t_0.

By cofinality there must be t_n in ρ such that $u \to^* t_n$.

Hence $t_n = u$ since u is a normal form.

Definition

A reduction $\rho = t_0 \to t_1 \to \dots$ is cofinal if for every $t_0 \to^* s$ there exists t_n in ρ such that $s \to^* t_n$.

Definition

A strategy $\mathcal{S}$ is cofinal if every every maximal $\mathcal{S}$ rewrite sequence is cofinal.

Theorem

Cofinal strategies are normalizing.

Proof.

Let $\mathcal{S}$ be a cofinal strategy.

Definition

A reduction $\rho = t_0 \to t_1 \to \dots$ is cofinal if for every $t_0 \to^* s$ there exists t_n in ρ such that $s \to^* t_n$.

Definition

A strategy $\mathcal{S}$ is cofinal if every every maximal $\mathcal{S}$ rewrite sequence is cofinal.

Theorem

Cofinal strategies are normalizing.

Proof.

Let $\mathcal{S}$ be a cofinal strategy. Let t_0 be a term that has a normal form u ($t_0 \to^* u$).

Definition

A reduction $\rho = t_0 \to t_1 \to \dots$ is cofinal if for every $t_0 \to^* s$ there exists t_n in ρ such that $s \to^* t_n$.

Definition

A strategy $\mathcal{S}$ is cofinal if every every maximal $\mathcal{S}$ rewrite sequence is cofinal.

Theorem

Cofinal strategies are normalizing.

Proof.

Let $\mathcal{S}$ be a cofinal strategy. Let t_0 be a term that has a normal form u ($t_0 \to^* u$).

Consider a maximal $\mathcal{S}$ rewrite sequence $\rho : t_0 \to t_1 \to \dots$ starting from t_0.

Definition

A reduction $\rho = t_0 \to t_1 \to \ldots$ is cofinal if for every $t_0 \to^* s$ there exists t_n in ρ such that $s \to^* t_n$.

Definition

A strategy $\mathcal{S}$ is cofinal if every every maximal $\mathcal{S}$ rewrite sequence is cofinal.

Theorem

Cofinal strategies are normalizing.

Proof.

Let $\mathcal{S}$ be a cofinal strategy. Let t_0 be a term that has a normal form u ($t_0 \to^* u$).

Consider a maximal $\mathcal{S}$ rewrite sequence $\rho : t_0 \to t_1 \to \ldots$ starting from t_0.

By cofinality there must be t_n in ρ such that $u \to^* t_n$.

Definition

A reduction $\rho = t_0 \to t_1 \to \ldots$ is cofinal if for every $t_0 \to^* s$ there exists t_n in ρ such that $s \to^* t_n$.

Definition

A strategy $\mathcal{S}$ is cofinal if every every maximal $\mathcal{S}$ rewrite sequence is cofinal.

Theorem

Cofinal strategies are normalizing.

Proof.

Let $\mathcal{S}$ be a cofinal strategy. Let t_0 be a term that has a normal form u ($t_0 \to^* u$).

Consider a maximal $\mathcal{S}$ rewrite sequence $\rho : t_0 \to t_1 \to \ldots$ starting from t_0.

By cofinality there must be t_n in ρ such that $u \to^* t_n$.

Hence $t_n = u$ since u is a normal form. ■

Theorem

For orthogonal TRSs, every fair strategy is cofinal.

Let $\tau : t_0 \to u_0$. We show that $u_0 \to^* t_n$ for some t_n in ρ.

$$\begin{array}{ccccccccccc} t_0 & \longrightarrow & t_1 & \longrightarrow & t_2 & \longrightarrow & \cdots & \longrightarrow & t_n & \longrightarrow & \cdots \\ \downarrow \tau & & \downarrow \tau/\rho_1 & & \downarrow \tau/\rho_2 & & & & \downarrow \varnothing & & \\ u_0 & \longrightarrow\!\!\!\!\!\rightarrow & u_1 & \longrightarrow\!\!\!\!\!\rightarrow & u_2 & \longrightarrow\!\!\!\!\!\rightarrow & \cdots & \longrightarrow\!\!\!\!\!\rightarrow & u_n & & \end{array}$$

Here ρ_i consists of the first i steps of ρ.

By fairness of ρ there exists n such that $\tau/\rho_n = \varnothing$. Hence $u_n = t_n$ and $u_0 \to^* t_n$.

The reduction $u_0 \to u_1 \to \ldots \to u_n \to t_{n+1} \to t_{n+2}$ is fair again.
(every redex occurrence in t_n is eventually secured)

By induction over the length of $t_0 \to^* u_0$ we get $u_0 \to^* t_n$ for some t_n in ρ. ∎

Theorem

For orthogonal TRSs, every fair strategy is cofinal.

Proof.

Let $\rho : t_0 \to t_1 \to \ldots$ be a fair rewrite sequence.

Theorem

For orthogonal TRSs, every fair strategy is cofinal.

Proof.

Let $\rho : t_0 \to t_1 \to \ldots$ be a fair rewrite sequence.

Let $\tau : t_0 \to u_0$. We show that $u_0 \to^* t_n$ for some t_n in ρ.

Theorem

For orthogonal TRSs, every fair strategy is cofinal.

Proof.

Let $\rho : t_0 \to t_1 \to \dots$ be a fair rewrite sequence.

Let $\tau : t_0 \to u_0$. We show that $u_0 \to^* t_n$ for some t_n in ρ.

$$\begin{array}{ccccccccccc}
t_0 & \longrightarrow & t_1 & \longrightarrow & t_2 & \longrightarrow & \cdots & \longrightarrow & t_n & \longrightarrow & \cdots \\
\downarrow \tau & & \downarrow \tau/\rho_1 & & \downarrow \tau/\rho_2 & & & & \downarrow \varnothing & & \\
u_0 & \longrightarrow\!\!\!\!\!\rightarrow & u_1 & \longrightarrow\!\!\!\!\!\rightarrow & u_2 & \longrightarrow\!\!\!\!\!\rightarrow & \cdots & \longrightarrow\!\!\!\!\!\rightarrow & u_n & &
\end{array}$$

Here ρ_i consists of the first i steps of ρ.

Theorem

For orthogonal TRSs, every fair strategy is cofinal.

Proof.

Let $\rho : t_0 \to t_1 \to \dots$ be a fair rewrite sequence.

Let $\tau : t_0 \to u_0$. We show that $u_0 \to^* t_n$ for some t_n in ρ.

$$\begin{array}{ccccccccccc} t_0 & \longrightarrow & t_1 & \longrightarrow & t_2 & \longrightarrow & \cdots & \longrightarrow & t_n & \longrightarrow & \cdots \\ \downarrow \tau & & \downarrow \tau/\rho_1 & & \downarrow \tau/\rho_2 & & & & \mid \varnothing & & \\ u_0 & \longrightarrow\!\!\!\!\!\rightarrow & u_1 & \longrightarrow\!\!\!\!\!\rightarrow & u_2 & \longrightarrow\!\!\!\!\!\rightarrow & \cdots & \longrightarrow\!\!\!\!\!\rightarrow & u_n & & \end{array}$$

Here ρ_i consists of the first i steps of ρ.

By fairness of ρ there exists n such that $\tau/\rho_n = \varnothing$.

Theorem

For orthogonal TRSs, every fair strategy is cofinal.

Proof.

Let $\rho : t_0 \to t_1 \to \dots$ be a fair rewrite sequence.

Let $\tau : t_0 \to u_0$. We show that $u_0 \to^* t_n$ for some t_n in ρ.

$$\begin{array}{ccccccccccc} t_0 & \longrightarrow & t_1 & \longrightarrow & t_2 & \longrightarrow & \cdots & \longrightarrow & t_n & \longrightarrow & \cdots \\ \downarrow \tau & & \downarrow \tau/\rho_1 & & \downarrow \tau/\rho_2 & & & & \mid \varnothing & & \\ u_0 & \longrightarrow\!\!\!\!\!\rightarrow & u_1 & \longrightarrow\!\!\!\!\!\rightarrow & u_2 & \longrightarrow\!\!\!\!\!\rightarrow & \cdots & \longrightarrow\!\!\!\!\!\rightarrow & u_n & & \end{array}$$

Here ρ_i consists of the first i steps of ρ.

By fairness of ρ there exists n such that $\tau/\rho_n = \varnothing$. Hence $u_n = t_n$ and $u_0 \to^* t_n$.

Theorem

For orthogonal TRSs, every fair strategy is cofinal.

Proof.

Let $\rho : t_0 \to t_1 \to \ldots$ be a fair rewrite sequence.

Let $\tau : t_0 \to u_0$. We show that $u_0 \to^* t_n$ for some t_n in ρ.

$$\begin{array}{ccccccccccc} t_0 & \longrightarrow & t_1 & \longrightarrow & t_2 & \longrightarrow & \cdots & \longrightarrow & t_n & \longrightarrow & \cdots \\ \downarrow \tau & & \downarrow \tau/\rho_1 & & \downarrow \tau/\rho_2 & & & & \downarrow \varnothing & & \\ u_0 & \longrightarrow\!\!\!\!\!\rightarrow & u_1 & \longrightarrow\!\!\!\!\!\rightarrow & u_2 & \longrightarrow\!\!\!\!\!\rightarrow & \cdots & \longrightarrow\!\!\!\!\!\rightarrow & u_n & & \end{array}$$

Here ρ_i consists of the first i steps of ρ.

By fairness of ρ there exists n such that $\tau/\rho_n = \varnothing$. Hence $u_n = t_n$ and $u_0 \to^* t_n$.

The reduction $u_0 \to u_1 \to \ldots \to u_n \to t_{n+1} \to t_{n+2}$ is fair again.
(every redex occurrence in t_n is eventually secured)

Theorem

For orthogonal TRSs, every fair strategy is cofinal.

Proof.

Let $\rho : t_0 \to t_1 \to \ldots$ be a fair rewrite sequence.

Let $\tau : t_0 \to u_0$. We show that $u_0 \to^* t_n$ for some t_n in ρ.

$$\begin{array}{ccccccccccc} t_0 & \longrightarrow & t_1 & \longrightarrow & t_2 & \longrightarrow & \cdots & \longrightarrow & t_n & \longrightarrow & \cdots \\ \downarrow \tau & & \downarrow \tau/\rho_1 & & \downarrow \tau/\rho_2 & & & & \downarrow \varnothing & & \\ u_0 & \longrightarrow\!\!\!\!\!\rightarrow & u_1 & \longrightarrow\!\!\!\!\!\rightarrow & u_2 & \longrightarrow\!\!\!\!\!\rightarrow & \cdots & \longrightarrow\!\!\!\!\!\rightarrow & u_n & & \end{array}$$

Here ρ_i consists of the first i steps of ρ.

By fairness of ρ there exists n such that $\tau/\rho_n = \varnothing$. Hence $u_n = t_n$ and $u_0 \to^* t_n$.

The reduction $u_0 \to u_1 \to \ldots \to u_n \to t_{n+1} \to t_{n+2}$ is fair again.
(every redex occurrence in t_n is eventually secured)

By induction over the length of $t_0 \to^* u_0$ we get $u_0 \to^* t_n$ for some t_n in ρ. ■

Definition

A TRS is left-normal if variables do not precede function symbols in left-hand sides (where the left-hand sides are written in prefix notation).

Definition

A TRS is left-normal if variables do not precede function symbols in left-hand sides (where the left-hand sides are written in prefix notation).

Example

- $\mathsf{f}(x,\ \mathsf{g}(y,z)) \to \mathsf{g}(y,\ \mathsf{f}(x,z))$

Definition

A TRS is left-normal if variables do not precede function symbols in left-hand sides (where the left-hand sides are written in prefix notation).

Example

- $\mathsf{f}(x,\ \mathsf{g}(y,z)) \to \mathsf{g}(y,\ \mathsf{f}(x,z))$ ☹

Definition

A TRS is left-normal if variables do not precede function symbols in left-hand sides (where the left-hand sides are written in prefix notation).

Example

- $\mathsf{f}(x,\ \mathsf{g}(y,z)) \to \mathsf{g}(y,\ \mathsf{f}(x,z))$ ☹
- $\mathsf{f}(\mathsf{g}(x,y),\ z) \to \mathsf{g}(x,\ \mathsf{g}(y,z))$

Definition

A TRS is left-normal if variables do not precede function symbols in left-hand sides (where the left-hand sides are written in prefix notation).

Example

- $\mathsf{f}(x,\ \mathsf{g}(y,z)) \to \mathsf{g}(y,\ \mathsf{f}(x,z))$ ☹
- $\mathsf{f}(\mathsf{g}(x,y),\ z) \to \mathsf{g}(x,\ \mathsf{g}(y,z))$ ☺

Definition

A TRS is left-normal if variables do not precede function symbols in left-hand sides (where the left-hand sides are written in prefix notation).

Example

- $\mathrm{f}(x,\ \mathrm{g}(y,z)) \to \mathrm{g}(y,\ \mathrm{f}(x,z))$ ☹
- $\mathrm{f}(\mathrm{g}(x,y),\ z) \to \mathrm{g}(x,\ \mathrm{g}(y,z))$ ☺

Theorem

Leftmost outermost strategy is normalizing for orthogonal left-normal TRSs.

Definition

A TRS is left-normal if variables do not precede function symbols in left-hand sides (where the left-hand sides are written in prefix notation).

Example

- $\mathsf{f}(x,\ \mathsf{g}(y,z)) \to \mathsf{g}(y,\ \mathsf{f}(x,z))$ ☹
- $\mathsf{f}(\mathsf{g}(x,y),\ z) \to \mathsf{g}(x,\ \mathsf{g}(y,z))$ ☺

Theorem

Leftmost outermost strategy is normalizing for orthogonal left-normal TRSs.

Remark

Combinatory Logic is left-normal

$$\mathsf{I}\,x \to x \qquad \mathsf{K}\,x\,y \to x \qquad \mathsf{S}\,x\,y\,z \to x\,z\,(y\,z)$$